

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
15 January 2004 (15.01.2004)

PCT

(10) International Publication Number  
WO 2004/005285 A1

(51) International Patent Classification<sup>7</sup>: C07D 453/02,  
487/08, A61K 31/439, 31/407, A61P 1/00, 11/00, 13/00 //  
(C07D 487/08, 209:00, 209:00)

(74) Agents: CRESSWELL, Thomas, Anthony et al.; J.A.  
Kemp & Co., 14 South Square, Gray's Inn, London WC1R  
5JJ (GB).

(21) International Application Number:  
PCT/EP2003/006708

(22) International Filing Date: 25 June 2003 (25.06.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
P 200201539 2 July 2002 (02.07.2002) ES

(71) Applicant (for all designated States except US): ALMI-  
RALE PRODEFARMA S.A. [ES/ES]; Ronda del Gen-  
eral Mitre, 151, E-08022 Barcelona (ES).

(72) Inventor; and

(75) Inventor/Applicant (for US only): PRAT QUINONES,  
Maria [ES/ES]; Calle Andrea Doria, 2, 1° 1ª, E-08003  
Barcelona (ES).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,  
MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC,  
SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG,  
US, UZ, VC, VN, YU, ZA, ZM, ZW.


(84) Designated States (regional): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,  
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,  
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,  
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: NEW QUINUCLIDINE AMIDE DERIVATIVES



(57) Abstract: New quinuclidine amide derivatives having the chemical structure of general formula (I) and pharmaceutically accept-  
able salts thereof including quaternary salts of formula (II) are disclosed; as well as processes for their preparation, pharmaceutical  
compositions 10 comprising them and their use in therapy as antagonists of M3 muscarinic receptors.

WO 2004/005285 A1

**QUINUCLIDINE AMIDE DERIVATIVES**

This invention relates to new therapeutically useful quinuclidine amide derivatives, to some processes for their preparation and to pharmaceutical compositions containing  
5 them.


The novel structures according to the invention are antimuscarinic agents with a potent and long lasting effect. In particular, these compounds show high affinity for M3 muscarinic receptors. This subtype of muscarinic receptor is present in glands and  
10 smooth muscle and mediates the excitatory effects of the parasympathetic system on glandular secretion and on the contraction of visceral smooth muscle (Chapter 6, Cholinergic Transmission, in H.P. Rang et al., Pharmacology, Churchill Livingstone, New York, 1995).

15 M3 antagonists are therefore known to be useful for treating diseases characterised by an increased parasympathetic tone, by excessive glandular secretion or by smooth muscle contraction (R.M. Eglen and S.S. Hegde, (1997), Drug News Perspect., 10(8):462-469).

Examples of this kind of diseases are respiratory disorders such as chronic obstructive  
20 pulmonary disease (COPD), bronchitis, bronchial hyperreactivity, asthma, cough and rhinitis; urological disorders such as urinary incontinence, pollakiuria, neurogenic or unstable bladder, cystospasm and chronic cystitis; gastrointestinal disorders such as irritable bowel syndrome, spastic colitis, diverticulitis and peptic ulceration; and cardiovascular disorders such as vagally induced sinus bradycardia (Chapter 7,  
25 Muscarinic Receptor Agonists and Antagonists, in Goodman and Gilman's The Pharmacological Basis of Therapeutics, 10th edition, McGraw Hill, New York, 2001).

The compounds of the invention can be used alone or in association with other drugs commonly regarded as effective in the treatment of these diseases. For example, they can  
30 be administered in combination with  $\beta_2$ -agonists, steroids, antiallergic drugs, phosphodiesterase IV inhibitors and/or leukotriene D4 (LTD4) antagonists for simultaneous, separate or sequential use in the treatment of a respiratory disease.

The new quinuclidine amide derivatives of the invention have the chemical structure of  
35 formula (I):




5 wherein


$R_1$  represents a hydrogen atom or a straight or branched, optionally substituted lower alkyl group;

10  $R_2$  represents a group of formula i) or ii)

i)



ii)



wherein

15  $R_3$  represents a group selected from phenyl, 2-furyl, 3-furyl, 2-thienyl or 3-thienyl;

$R_4$  represents a group selected from optionally substituted lower alkyl, optionally substituted lower alkenyl, optionally substituted lower alkynyl, cycloalkyl, cycloalkylmethyl, phenyl, benzyl, phenethyl, 2-furyl, 3-furyl, 2-thienyl or 3-thienyl;

20

and  $R_5$  represents a hydrogen atom or a hydroxy, methyl, or  $-CH_2OH$  group;

the benzene rings in formula ii) and the cyclic groups represented by  $R_3$  and  $R_4$  being each independently optionally substituted by one, two or three substituents selected from

25 halogen, straight or branched, optionally substituted lower alkyl, hydroxy, straight or branched, optionally substituted lower alkoxy, nitro, cyano,  $-CO_2R'$  or  $-NR'R''$ , wherein  $R'$

and R'' each independently represents a hydrogen atom or a straight or branched, optionally substituted lower alkyl group or R' and R'' together with the atom to which they are attached form a cyclic group;

5 Q represents a single bond or a -CH<sub>2</sub>-, -CH<sub>2</sub>-CH<sub>2</sub>-, -O-, -O-CH<sub>2</sub>-, -S-, -S-CH<sub>2</sub>- or -CH=CH- group;

p is 1 or 2 and the amide group is at positions 2, 3 or 4 of the azabicyclic ring;

10 or pharmaceutically acceptable salts thereof; including quaternary ammonium salts;

and all individual stereoisomers and mixtures thereof;


with the proviso that when p is 2, the amide moiety is in position 3 of the quinuclidine ring,

15 R<sub>1</sub> is hydrogen and R<sub>3</sub> and R<sub>4</sub> are both unsubstituted phenyl, then

-when said compound is not a pharmaceutically acceptable salt or is a HCl salt, then R<sub>5</sub> cannot be one of hydrogen or hydroxy; and

- when said compound is a quaternary ammonium salt having a methyl group attached to  
20 the nitrogen atom of the quinuclidine ring, then R<sub>5</sub> cannot be hydroxy.

The compounds of the invention include quaternary ammonium salts of formula (II).



(II)

25 wherein R<sub>1</sub>, R<sub>2</sub> and p are as defined above;

m is an integer from 0 to 8;

n is an integer from 0 to 4:

A represents a group selected from  $-\text{CH}_2-$ ,  $-\text{CH}=\text{CR}'-$ ,  $-\text{CR}'=\text{CH}-$ ,  $-\text{CR}'\text{R}''-$ ,  $-\text{C}(\text{O})-$ ,  $-\text{O}-$ ,  $-\text{S}-$ ,  $-\text{S}(\text{O})-$ ,  $-\text{S}(\text{O})_2-$  and  $-\text{NR}'-$ , wherein  $\text{R}'$  and  $\text{R}''$  are as defined above;

$\text{R}_6$  represents a hydrogen atom, or a group selected from straight or branched, optionally substituted lower alkyl, hydroxy, straight or branched, optionally substituted lower alkoxy, cyano, nitro,  $-\text{CH}=\text{CR}'\text{R}''$ ,  $-\text{C}(\text{O})\text{OR}'$ ,  $-\text{OC}(\text{O})\text{R}'$ ,  $-\text{SC}(\text{O})\text{R}'$ ,  $-\text{C}(\text{O})\text{NR}'\text{R}''$ ,  $-\text{NR}'\text{C}(\text{O})\text{OR}''$ ,  $-\text{NR}'\text{C}(\text{O})\text{NR}''$ , cycloalkyl, phenyl, naphthanelyl, 5,6,7,8-tetrahydronaphthanelyl, benzo[1,3]dioxolyl, heteroaryl or heterocyclyl;  $\text{R}'$  and  $\text{R}''$  being as defined above; and wherein the cyclic groups represented by  $\text{R}_6$  are optionally substituted by one, two or three substituents selected from halogen, hydroxy, straight or branched, optionally substituted lower alkyl, phenyl,  $-\text{OR}'$ ,  $-\text{SR}'$ ,  $-\text{NR}'\text{R}''$ ,  $-\text{NHCOR}'$ ,  $-\text{CONR}'\text{R}''$ ,  $-\text{CN}$ ,  $-\text{NO}_2$  and  $-\text{COOR}'$ ;  $\text{R}'$  and  $\text{R}''$  being as defined above;

$\text{X}^-$  represents a pharmaceutically acceptable anion of a mono or polyvalent acid.

and all individual stereoisomers and mixtures thereof;

with the proviso that when  $p$  is 2, the amide moiety is in position 3 of the quinuclidine ring,  $\text{R}_1$  is hydrogen,  $\text{R}_3$  and  $\text{R}_4$  are both unsubstituted phenyl and  $\text{R}_5$  is hydroxy, then in the compounds of formula (II) the sequence  $\text{R}_6 - (\text{CH}_2)_n - \text{A} - (\text{CH}_2)_m -$  cannot be a methyl group.

Further objectives of the present invention are to provide processes for preparing said compounds; pharmaceutical compositions comprising an effective amount of said compounds; the use of the compounds in the manufacture of a medicament for the treatment of diseases susceptible of being improved by antagonism of M3 muscarinic receptors; and methods of treatment of diseases susceptible to amelioration by antagonism of M3 muscarinic receptors, which methods comprise the administration of the compounds of the invention to a subject in need of treatment.

Certain quinuclidine amide derivatives, which fall outside the scope of the present invention, have been disclosed in JP 09328469 and WO 9316048.

As used herein, an alkyl group or moiety can be straight or branched, and is typically a lower alkyl group or moiety. A lower alkyl group or moiety contains 1 to 8, preferably 1 to 6

and more preferably 1 to 4 carbon atoms. In particular it is preferred that such an alkyl group or moiety is represented by a methyl, ethyl, propyl, including i-propyl or butyl, including n-butyl, sec-butyl and tert-butyl group or moiety.

- 5 As used herein, an alkenyl group or moiety can be straight or branched, and is typically a lower alkenyl group or moiety. A lower alkenyl group or moiety contains 2 to 8, preferably 2 to 6 and more preferably 2 to 4 carbon atoms. In particular it is preferred that such an alkenyl group or moiety is represented by a vinyl, allyl or 1-propenyl group or moiety.
- 10 As used herein, an alkynyl group or moiety can be straight or branched, and is typically a lower alkynyl group or moiety. A lower alkynyl group or moiety contains 2 to 8, preferably 2 to 6 and more preferably 2 to 4 carbon atoms. In particular it is preferred that such an alkynyl group or moiety is represented by a 3-butylnyl or 1-propynyl group or moiety.
- 15 Optionally substituted lower alkyl, alkenyl or alkynyl groups mentioned herein include straight or branched lower alkyl, alkenyl or alkynyl groups as defined above, which may be unsubstituted or substituted in any position by one or more substituents, for example by 1, 2 or 3 substituents. When two or more substituents are present, each substituent may be the same or different. The substituent(s) are typically halogen atoms, preferably fluoride
- 20 atoms, and hydroxy or alkoxy groups.

- Alkoxy groups mentioned herein are typically lower alkoxy groups, that is groups containing from 1 to 8, preferably 1 to 6 and more preferably 1 to 4 carbon atoms, the hydrocarbon chain being branched or straight and optionally substituted in any position by
- 25 one or more substituents, for example by 1, 2 or 3 substituents. When two or more substituents are present, each substituent may be the same or different. The substituent(s) are typically halogen atoms, most preferably fluoride atoms, and hydroxy groups. Preferred optionally substituted alkoxy groups include methoxy, ethoxy, n-propoxy, i-propoxy, n-butoxy, sec-butoxy, t-butoxy, trifluoromethoxy and difluoromethoxy,
- 30 hydroxymethoxy, hydroxyethoxy or 2-hydroxypropoxy.

Cyclic groups mentioned herein include carbocyclic and heterocyclic groups. The cyclic groups can contain one or more rings. Carbocyclic groups may be aromatic or alicyclic, for example cycloalkyl groups. Heterocyclic groups also include heteroaryl groups.

Cycloalkyl groups and alicyclic groups mentioned herein, unless otherwise specified, typically contain from 3 to 7 carbon atoms. Cycloalkyl groups and alicyclic rings of 3 to 7 carbon atoms include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl.

- 5 As used herein an aromatic group typically contains from 5 to 14, preferably 5 to 10 carbon atoms. Examples of aromatic groups include phenyl and naphthalenyl.

- A heterocyclic or heteroaromatic group mentioned herein is typically a 5 to 10 membered group, such as a 5, 6 or 7 membered group, containing one or more heteroatoms selected from N, S and O. Typically, 1, 2, 3 or 4 heteroatoms are present, preferably 1 or 2 heteroatoms. A heterocyclic or heteroaromatic group may be a single ring or two or more fused rings wherein at least one ring contains a heteroatom. Examples of heterocyclic groups include piperidyl, pyrrolidyl, piperazinyl, morpholinyl, thiomorpholinyl, pyrrolyl, imidazolyl, imidazolidinyl, pyrazolyl, indolyl, isoindolyl, pyridyl, pyrazinyl, pyrimidinyl, 10 pyridazinyl, indolizyl, isoindolyl, indolyl, indazolyl, purinyl, quinolizyl, isoquinolyl, quinolyl, phthalazinyl, naphthyridinyl, quinoxalinyl, quinazolinyl, cinnolyl, pteridinyl, quinuclidinyl, triazolyl, pyrazolyl, tetrazolyl, thienyl and dioxolyl. Examples of heteroaromatic groups include pyridyl, thienyl, furyl, pyrrolyl, imidazolyl, benzothiazolyl, pyridinyl, pyrazolyl, pyrazinyl, pyrimidinyl, pyridazinyl, indolyl, indazolyl, purinyl, quinolyl, 20 isoquinolyl, phthalazinyl, naphthyridinyl, quinoxalinyl, quinazolinyl, cinnolyl, triazolyl and pyrazolyl.

- As used herein a halogen atom includes a fluorine, chlorine, bromine or iodine atom, typically a fluorine, chlorine or bromine atom.

25

- As used herein, a pharmaceutically acceptable salt is a salt with a pharmaceutically acceptable acid or base. Pharmaceutically acceptable acids include both inorganic acids, for example hydrochloric, sulphuric, phosphoric, diphosphoric, hydrobromic and nitric acid and organic acids, for example citric, fumaric, formic, maleic, malic, ascorbic, succinic, 30 tartaric, benzoic, acetic, methanesulphonic, ethanesulphonic, benzenesulphonic or p-toluenesulphonic acid. Pharmaceutically acceptable bases include alkali metal (e.g. sodium or potassium) and alkali earth metal (e.g. calcium or magnesium) hydroxides and organic bases, for example alkyl amines, aralkyl amines and heterocyclic amines. Preferred salts of the compounds of formula (I) are those formed between the nitrogen 35 atom of the azabicyclic ring and an inorganic or an organic acid

Other preferred salts according to the invention are quaternary ammonium compounds of formula (II), wherein an equivalent of an anion ( $X^-$ ) is associated with the positive charge of the N atom.  $X^-$  may be an anion of various mineral acids such as, for example, chloride, bromide, iodide, sulphate, nitrate, phosphate, or an anion of an organic acid such as, for example, acetate, formate, maleate, fumarate, citrate, oxalate, succinate, tartrate, malate, mandelate, trifluoroacetate, methanesulphonate and p-toluenesulphonate.  $X^-$  is preferably an anion selected from chloride, bromide, iodide, sulphate, nitrate, acetate, maleate, oxalate, succinate or trifluoroacetate. More preferably  $X^-$  is chloride, bromide, trifluoroacetate or methanesulphonate.

Preferred compounds of formulae (I) or (II) are those wherein  $R_1$  is hydrogen, methyl or ethyl, most preferably hydrogen.

In some preferred embodiments of the invention  $R_2$  is a group of formula i), wherein  $R_3$  is a group, which is optionally substituted with one or more halogen atom(s), selected from phenyl, 2-thienyl, 3-thienyl or 2-furyl. More preferably  $R_2$  is a group of formula i), wherein  $R_3$  represents a group phenyl, 2-thienyl or 2-furyl which are optionally substituted with one or more halogen atom(s). Most preferably,  $R_3$  is phenyl or 2-thienyl,

20

Further preferred compounds having a group of formula i) are those wherein  $R_4$  represents a linear group selected from ethyl, n-butyl, vinyl, allyl, 1-propenyl and 1-propynyl, or a group, which is optionally substituted with one or more halogen atom(s), methyl or methoxy group(s), selected from cyclopentyl, cyclohexyl, phenyl, benzyl, phenethyl, 2-thienyl and 3-furyl. More preferably compounds having a group of formula i) are those wherein  $R_4$  represents a linear group selected from ethyl, n-butyl, vinyl, allyl and 1-propynyl, or a group, which is optionally substituted with one or more halogen atom(s), methyl or methoxy group(s), selected from cyclopentyl, phenyl, benzyl, phenethyl and 2-thienyl. Most preferably,  $R_4$  is ethyl, n-butyl, vinyl, allyl, cyclopentyl, phenyl, benzyl or 2-thienyl.

30

In other preferred embodiments of the invention  $R_2$  is a group of formula ii), wherein Q represents a single bond or an oxygen atom.



In the most preferred embodiments of the invention  $R_5$  in groups i) or ii) is a hydrogen atom or a hydroxy group.

Preferably, in the compounds of formulae (I) or (II)  $p$  is 2 and the amide group is at positions 3 or 4 of the azabicyclic ring, most preferably at position 3.

Particularly preferred compounds of formula (II) are those wherein  $m$  is an integer from 0 to 6 and  $n$  is an integer from 0 to 4; most preferably  $m$  is 0 to 5 and  $n$  0 to 2;  $A$  represents a group selected from  $-CH_2-$ ,  $-CH=CH-$ ,  $-O-$ ,  $-C(O)-$ ,  $-NR'-$ , and  $-S-$ ; most preferably  $A$  is  $-CH_2-$ ,  $-CH=CH-$ ,  $-O-$ ; and  $R_6$  is a hydrogen atom, a cyano group, a nitro group, a  $-C(O)OR'$  group, a  $-OC(O)R'$  group, a  $-SC(O)R'$  group, a  $-CH=CH_2$  group, a  $-CH=CR'R''$  group, a  $C(O)NR'R''$  group, a straight or branched  $C_1-C_4$  alkyl group, which is optionally substituted with one or more halogen atom(s), a straight  $C_1-C_4$  alkoxy group, which is optionally substituted with one or more halogen atom(s) or hydroxy group(s), or a cyclic group, which is optionally substituted with one or more substituents selected from halogen atoms, groups of formula  $-C(O)NR'R''$  or methyl, hydroxy, nitro or phenyl groups, the cyclic group being selected from cyclohexyl, phenyl, 5,6,7,8-tetrahydronaphthanelyl, 2-thienyl, 1-pyrrolidinyl, 1-pyrrolyl, benzo[1,3]dioxolyl, 2-benzothiazolyl, naphthalenyl and dioxolyl. More preferably compounds of formula (II) are those wherein  $m$  is an integer from 0 to 5 and  $n$  is an integer from 0 to 2;  $A$  represents a group selected from  $-CH_2-$ ,  $-CH=CH-$ ,  $-O-$ ,  $-C(O)-$ ,  $-NR'-$ , and  $-S-$ ; and  $R_6$  is a hydrogen atom, a cyano group, a  $-C(O)OR'$  group, a  $-OC(O)R'$  group, a  $-SC(O)R'$  group, a  $-CH=CH_2$  group, a  $-C(O)NR'R''$  group, a straight or branched  $C_1-C_4$  alkyl group, a trifluoromethyl, or a cyclic group selected from cyclohexyl, 5,6,7,8-tetrahydronaphthanelyl, 2-thienyl, 1-pyrrolyl, benzo[1,3]dioxolyl, 2-benzothiazolyl, naphthalenyl, dioxolyl and phenyl, which is optionally substituted with one or more substituents selected from halogen atoms, groups of formula  $-C(O)NR'R''$ , methyl, hydroxy and phenyl groups. Most preferably  $R_6$  is selected from hydrogen, straight  $C_1-C_4$  alkyl,  $-CH=CH_2$ , cyclohexyl, phenyl which is unsubstituted or substituted with one or two substituents selected from methyl groups and hydroxy groups, 5,6,7,8-tetrahydronaphthanelyl and 2-thienyl.

Most preferred are compounds of formula (II) wherein the sequence  $R_6 - (CH_2)_n - A - (CH_2)_m -$  is one of methyl, 3-phenoxypropyl, 3-(3-hydroxyphenoxy)propyl, allyl, heptyl, 3-phenylpropyl, 3-phenylallyl, 2-phenoxyethyl, 2-benzyloxyethyl, cyclohexylmethyl, 3-

(5,6,7,8-tetrahydronaphthalen-2-yloxy)propyl, 5-(2,6-dimethylphenoxy)pentyl, 3-thien-2-ylpropyl or 3-cyclohexylpropyl and X<sup>-</sup> is bromide or trifluoroacetate.

The compounds of the present invention represented by formulae (I) or (II) described  
 5 above may have one or more asymmetric atoms, e.g. the carbon in position 3 of the quinuclidine ring; the carbon substituted by R<sub>3</sub>, R<sub>4</sub> and R<sub>5</sub> in the compounds wherein R<sub>2</sub> is a group of formula i); or the carbon which is attached to the carbonyl group in the compounds wherein R<sub>2</sub> is a group of formula ii). Each of these asymmetric atoms may have R- or S-configuration. The single isomers and mixtures of the isomers fall within the  
 10 scope of the invention.

Particular individual compounds of formula (I) include:

- N-(1-Azabicyclo[2.2.2]oct-3-yl)-2-hydroxy-2,2-dithien-2-ylacetamide
- 15 N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2,2-dithien-2-ylacetamide
- N-[(3S)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2,2-dithien-2-ylacetamide
- N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2,2-dithien-2-ylacetamide
- N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-cyclopentyl-2-hydroxy-2-thien-2-ylacetamide
- N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2-thien-2-ylpent-4-enamide
- 20 (2\*)-N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2-thien-2-ylbutanamide  
 (diastereomer 1)
- (2\*)-N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2-thien-2-ylbutanamide  
 (diastereomer 2)
- (2\*)-N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2-thien-2-ylbut-3-enamide  
 25 (diastereomer 1)
- (2\*)-N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2-thien-2-ylbut-3-enamide  
 (diastereomer 2)
- N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2,3-diphenylpropanamide
- N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-cyclopentyl-2-hydroxy-2-phenylacetamide
- 30 N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-phenylhexanamide
- N-(1-Azabicyclo[2.2.2]oct-3-yl)-9H-xanthene-9-carboxamide
- N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-9H-xanthene-9-carboxamide
- N-[(3S)-1-Azabicyclo[2.2.2]oct-3-yl]-9H-xanthene-9-carboxamide
- N-(1-Azabicyclo[2.2.2]oct-3-yl)-9-hydroxy-9H-fluorene-9-carboxamide
- 35 N-1-Azabicyclo[2.2.2]oct-3-yl-N-methyl-9H-xanthene-9-carboxamide

N-1-Azabicyclo[2.2.2]oct-3-yl-N-ethyl-9H-xanthene-9-carboxamide

N-1-Azabicyclo[2.2.2]oct-4-yl-2-cyclopentyl-2-hydroxy-2-thien-2-ylacetamide

N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2-phenyl-2-thien-2-ylacetamide

N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-(5-bromothien-2-yl)-2-(4-fluoro-3-methylphenyl)-2-hydroxyacetamide

N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-fur-2-yl-2-hydroxypent-3-ynamide

N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-fur-2-yl-2-hydroxy-4-(4-methoxyphenyl)butanamide

N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-fur-3-yl-2-hydroxy-2-thien-2-ylacetamide

N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-cyclohexyl-2-hydroxy-2-thien-3-ylacetamide

10 N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-9-hydroxy-9H-fluorene-9-carboxamide

N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-N-methyl-9H-xanthene-9-carboxamide

(2S)-N-[(3R)-1-azabicyclo[2.2.2]oct-3-yl]-2-cyclopentyl-2-hydroxy-2-thien-2-ylacetamide

15 ((\*) Configuration not assigned; either the (2R)- or the (2S)- isomers of the above compounds may be produced).

Particular individual compounds of formula (II) include:

3-(2-Hydroxy-2,2-dithien-2-ylacetyl-amino)-1-methyl-1-azoniabicyclo[2.2.2]octane bromide

20 3-(2-Hydroxy-2,2-dithien-2-ylacetyl-amino)-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane bromide

(3R)-3-(2-Hydroxy-2,2-dithien-2-ylacetyl-amino)-1-methyl-1-azoniabicyclo[2.2.2]octane trifluoroacetate

25 (3R)-1-Allyl-3-(2-hydroxy-2,2-dithien-2-ylacetyl-amino)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

(3R)-1-Heptyl-3-(2-hydroxy-2,2-dithien-2-ylacetyl-amino)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

(3R)-3-(2-Hydroxy-2,2-dithien-2-ylacetyl-amino)-1-(3-phenylpropyl)-1-azoniabicyclo[2.2.2]octane bromide

30 (3R)-3-(2-Hydroxy-2,2-dithien-2-ylacetyl-amino)-1-((E)-3-phenylallyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

(3R)-3-(2-Hydroxy-2,2-dithien-2-ylacetyl-amino)-1-(2-phenoxyethyl)-1-azoniabicyclo[2.2.2]octane bromide

35 (3R)-3-(2-Hydroxy-2,2-dithien-2-ylacetyl-amino)-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane bromide

- (3R)-3-(2-Hydroxy-2,2-dithien-2-ylacetyl amino)-1-[3-(3-hydroxyphenoxy)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-1-(2-Benzoyloxyethyl)-3-(2-hydroxy-2,2-dithien-2-ylacetyl amino)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 5 (3R)-3-(2-Hydroxy-2,2-dithien-2-ylacetyl amino)-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane bromide
- (3S)-3-(2-Hydroxy-2,2-dithien-2-ylacetyl amino)-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane bromide
- (3R)-3-(2,2-Dithien-2-ylacetyl amino)-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane
- 10 bromide
- 1-Methyl-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane bromide
- 1-(3-Phenoxypropyl)-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane bromide
- (3R)-1-(3-Phenoxypropyl)-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane bromide
- 15 (3S)-1-Allyl-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3S)-1-Heptyl-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 20 (3S)-1-Cyclohexylmethyl-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3S)-1-(3-Cyclohexylpropyl)-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3S)-1-(3-Phenoxypropyl)-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane bromide
- 25 (3S)-1-[3-(5,6,7,8-Tetrahydronaphthalen-2-yloxy)propyl]-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3S)-1-[5-(2,6-Dimethylphenoxy)pentyl]-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 30 3-[(9-Hydroxy-9H-fluoren-9-yl)carbonyl]amino-1-methyl-1-azoniabicyclo[2.2.2]octane bromide
- 3-[(9-Hydroxy-9H-fluoren-9-yl)carbonyl]amino-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane bromide
- (3R)-1-[3-(2-Carbamoylphenoxy)propyl]-3-[(9-Hydroxy-9H-fluoren-9-yl)carbonyl]amino-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 35

- (3R)-1-[4-(4-Fluorophenyl)-4-oxobutyl]-3-[[9-Hydroxy-9H-fluoren-9-yl]carbonyl]amino]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[[9-Hydroxy-9H-fluoren-9-yl]carbonyl]amino]-1-(4-oxo-4-thien-2-ylbutyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 5 (3R)-3-[[9-Hydroxy-9H-fluoren-9-yl]carbonyl]amino]-1-[3-(methylphenylamino)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[[9-Hydroxy-9H-fluoren-9-yl]carbonyl]amino]-1-(3-phenylsulfanylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 3-[Methyl-(9H-xanthen-9-ylcarbonyl)amino]-1-(3-pyrrol-1-ylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 10 1-[3-(Biphenyl-4-yloxy)propyl]-3-[ethyl-(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-1-[3-(Benzo[1,3]dioxol-5-yloxy)propyl]-3-(2-hydroxy-2-phenyl-2-thien-2-ylacetyl-amino)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 15 (3R)-1-[3-(Benzothiazol-2-yloxy)propyl]-3-(2-hydroxy-2-phenyl-2-thien-2-ylacetyl-amino)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-(2-Fur-2-yl-2-hydroxypent-3-ynoylamino)-1-[3-(naphthalen-1-yloxy)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[2-Fur-2-yl-2-hydroxy-4-(4-methoxyphenyl)butyrylamino]-1-[6-(4-phenylbutoxy)hexyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 20 (3R)-3-(2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetyl-amino)-1-(2-hydroxyethyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-(2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetyl-amino)-1-(2-ethoxyethyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 25 (3R)-3-(2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetyl-amino)-1-[2-[2-(2-hydroxyethoxy)ethoxy]ethyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-(2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetyl-amino)-1-(4,4,4-trifluorobutyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-1-(4-Acetoxybutyl)-3-[2-(5-bromothien-2-yl)-2-(4-fluoro-3-methylphenyl)-2-hydroxyacetyl-amino]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 30 (3R)-3-[2-(5-Bromothien-2-yl)-2-(4-fluoro-3-methylphenyl)-2-hydroxyacetyl-amino]-1-(4-ethoxycarbonylbutyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-1-(3-Acetylsulfanylpropyl)-3-[2-(5-bromothien-2-yl)-2-(4-fluoro-3-methylphenyl)-2-hydroxyacetyl-amino]-1-azoniabicyclo[2.2.2]octane trifluoroacetate

(3R)-1-(3-Cyanopropyl)-3-(2-hydroxy-2-phenyl-2-thien-2-ylacetyl-amino)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

(3R)-1-(2-Carbamoylethyl)-3-(2-hydroxy-2-thien-2-ylpent-4-enoylamino)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

5 (3R)-1-(2-[1,3]Dioxolan-2-ylethyl)-3-(2-hydroxy-2-thien-2-ylpent-4-enoylamino)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

4-(2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetyl-amino)-1-(4-methylpent-3-enyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

(3R)-1-[3-(2-Carbamoylphenoxy)propyl]-3-[[ (9-Hydroxy-9H-fluoren-9-yl)carbonyl]amino]-1-azoniabicyclo[2.2.2]octane formate

(3R)-1-[4-(4-Fluorophenyl)-4-oxobutyl]-3-[[ (9-Hydroxy-9H-fluoren-9-yl)carbonyl]amino]-1-azoniabicyclo[2.2.2]octane formate.

(3R)-3-[[ (9-Hydroxy-9H-fluoren-9-yl)carbonyl]amino]-1-[3-(methylphenylamino)propyl]-1-azoniabicyclo[2.2.2]octane chloride

15 (3R)-3-[[ (9-Hydroxy-9H-fluoren-9-yl)carbonyl]amino]-1-(3-phenylsulfanylpropyl)-1-azoniabicyclo[2.2.2]octane formate

(3R)-3-[Methyl-(9H-xanthen-9-ylcarbonyl)amino]-1-(3-pyrrol-1-ylpropyl)-1-azoniabicyclo[2.2.2]octane bromide

(3R)-1-[3-(Biphenyl-4-yloxy)propyl]-3-[methyl-(9H-xanthene-9-carbonyl)amino]-1-azoniabicyclo[2.2.2]octane chloride

20 (3R)-3-(2-Fur-2-yl-2-hydroxypent-3-ynoylamino)-1-[3-(naphthalen-1-yloxy)propyl]-1-azoniabicyclo[2.2.2]octane chloride

(3R)-1-[3-(Benzo[1,3]dioxol-5-yloxy)propyl]-3-(2-fur-2-yl-2-hydroxypent-3-ynoylamino)-1-azonia-bicyclo[2.2.2]octane bromide

25 (3R)-1-[3-(Benzothiazol-2-yloxy)propyl]-3-(2-fur-2-yl-2-hydroxypent-3-ynoylamino)-1-azonia-bicyclo[2.2.2]octane chloride

(3R)-3-[[ (2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetyl]amino]-1-(2-hydroxyethyl)-1-azoniabicyclo[2.2.2]octane bromide

(3R)-3-[[ (2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetyl]amino]-1-(2-ethoxyethyl)-1-azoniabicyclo[2.2.2]octane formate


30 (3R)-3-[[ (2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetyl]amino]-1-(4,4,4-trifluorobutyl)-1-azoniabicyclo[2.2.2]octane bromide

(3R)-1-(4-Acetoxybutyl)-3-[2-(5-bromothien-2-yl)-2-(4-fluoro-3-methylphenyl)-2-hydroxyacetyl-amino]-1-azoniabicyclo[2.2.2]octane bromide

- (3R)-3-[2-(5-Bromothien-2-yl)-2-(4-fluoro-3-methylphenyl)-2-hydroxyacetyl-amino]-1-(4-ethoxycarbonylbutyl)-1-azoniabicyclo[2.2.2]octane bromide
- (3R)-1-(3-Acetylsulfanylpropyl)-3-[2-(5-bromothien-2-yl)-2-(4-fluoro-3-methylphenyl)-2-hydroxyacetyl-amino]-1-azoniabicyclo[2.2.2]octane formate
- 5 (3R)-1-(3-Cyanopropyl)-3-[2-fur-2-yl-2-hydroxy-4-(4-methoxyphenyl)butyrylamino]-1-azoniabicyclo[2.2.2]octane bromide
- (3R)-1-(2-Carbamoylethyl)-3-[2-fur-2-yl-2-hydroxy-4-(4-methoxyphenyl)butyrylamino]-1-azoniabicyclo[2.2.2]octane formate
- (3R)-1-(2-[1,3]Dioxolan-2-yl-ethyl)-3-[2-fur-2-yl-2-hydroxy-4-(4-methoxyphenyl)butyrylamino]-1-azoniabicyclo[2.2.2]octane bromide
- 10

In accordance with another embodiment, the present invention provides processes for preparing the novel quinuclidine amide derivatives of formulae (I) and (II)

- 15 The compounds of formula (I) may be prepared from the corresponding carboxylic acid of formula (III) following two different methods (a) and (b), illustrated in the following scheme.




20

Some compounds of formula (IV) are commercially available, for example 3-aminoquinuclidine (Aldrich, dihydrochloride), (3R)-aminoquinuclidine (Aldrich, Finorga, dihydrochloride) and (3S)-aminoquinuclidine (Aldrich, dihydrochloride).

(3R)-N-methylquinuclidin-3-amine has been prepared as described in the experimental section.

- 5 Those compounds of formula (I), wherein  $R_2$  is a group of formula i) and  $R_5$  is an hydroxy group may also be prepared from the compounds of formula (V) by reaction with the corresponding organometallic derivative.



10

The compounds of formula (V) may be prepared from the corresponding glyoxylic acids following methods equivalent to methods (a) and (b) described in Scheme 1

15 Certain compounds of formula (V) are novel and fall within the scope of the present invention. In particular:

N-1-azabicyclo[2.2.2]oct-3-yl-2-oxo-2-thien-2-ylacetamide

N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-oxo-2-thien-2-ylacetamide


N-[(3S)-1-Azabicyclo[2.2.2]oct-3-yl]-2-oxo-2-thien-2-ylacetamide

20

The quaternary ammonium derivatives of formula (II) may be prepared, as illustrated in the following scheme, by reaction of the compounds of formula (I) with an alkylating agent of formula (VI) using two possible methods -(d) or (e)-, described in detail in the experimental section. Method (e) involves the use of solid phase extraction techniques


25 that allow the parallel preparation of several compounds.





In formula (VI), W represents any suitable leaving group, preferably a group X<sup>-</sup> as defined above for the compounds of formula (II). When W is a leaving group other than X<sup>-</sup>, the quaternary ammonium salt of formula (II) is produced from the product of method (d) or (e) by an exchange reaction according to standard methods to replace the anion W<sup>-</sup> with the desired anion X<sup>-</sup>.

Those compounds of general formula (VI) which are not commercially available have been prepared according to standard methods. For example, compounds wherein n=0 and A = -O-, -S-, NR' were obtained by reaction of the corresponding alcohol, thiol or amine derivative, or its sodium or potassium salt, with an alkylating agent of general formula Y-(CH<sub>2</sub>)<sub>m</sub>-W, wherein W is as defined above; most preferably W is a halogen atom and Y a halogen atom or a sulphonate ester. In other examples, compounds of general formula (VI) wherein n is at least 1 were synthesised from the corresponding alcohol derivative of general formula (VII) by methods known in the art.



The compounds of formulae (I) and (II) can have one or more asymmetric carbons. All possible stereoisomers, single isomers and mixtures of isomers are also included within

the scope of the present invention. The diastereomers of the compounds may be separated by conventional methods, for example by chromatography or crystallisation.

The structures of the compounds were confirmed by <sup>1</sup>H-NMR and MS. The NMR spectra were recorded using either a Varian 300 MHz instrument or a Bruker DPX-250 instrument. Chemical shifts are expressed as parts per million (δ) from the internal reference tetramethylsilane. The purity of the compounds was determined by HPLC, using reverse phase chromatography on a Waters instrument. Molecular ions were produced by electrospray ionisation mass spectrometry on a Hewlett Packard instrument.

#### Method (a)

##### **Preparation of N-[(3R)-1-azabicyclo[2.2.2]oct-3-yl]-2-phenylhexanamide (Example 27)**

2-phenylhexanoic acid was prepared by alkylation of phenylacetic acid with 1-chlorobutane following a standard method. Oxalyl chloride (0.88 ml, 0.0101 mol) was then added to a solution of 1.62 g of 2-phenylhexanoic acid (0.0084 mol) and dimethylformamide (DMF, one drop) in 25 ml of CHCl<sub>3</sub> (ethanol free) at 0°C. The mixture was stirred and allowed to warm at room temperature. After an hour at this temperature the solvents were evaporated and the residue was dissolved in CHCl<sub>3</sub> and evaporated again. This procedure was repeated two times. The oil obtained was dissolved in CHCl<sub>3</sub> and the solution obtained cooled at 0°C. A solution of (3R)-aminoquinuclidine (1.28 g, 0.0101 mol) in CHCl<sub>3</sub> was added. The mixture was stirred and allowed to warm at room temperature. After 1 hour at this temperature, the reaction mixture was diluted with CHCl<sub>3</sub> and washed with a 10% solution of aqueous potassium carbonate, then washed with water, dried over Na<sub>2</sub>SO<sub>4</sub> and evaporated to give 3.57 g of an oil which was purified by column chromatography (silica gel, CHCl<sub>3</sub>:MeOH:NH<sub>4</sub>OH 90:10:1 as eluent). Appropriate fractions were combined and evaporated to give 2.42 g of a solid that after treatment with isopropyl ether yielded 1.59 g (63.1%) of the title product as a mixture of diastereomers.

mp: 136°C.

MS [M+1]<sup>+</sup>: 301

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): (mixture of diastereomers 50:50) δ 0.84 (t, 3H), 1.10-1.60 (m, 9H), 1.70 (m, 1H), 1.90 (m, 1H), 2.25-2.50 (m, 1H), 2.50-2.80 (m, 4H), 2.94 & 3.05 (m, 1H), 3.50 (m, 1H), 3.65 (m, 1H), 7.15-7.40 (m, 5H), 8.04 (m, 1H, NH).

**Preparation of N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-N-methyl-9H-xanthene-9-carboxamide (Example 47)**

Oxalyl chloride (0.540 ml, 0.0062 mol) was added to a solution of 9H-xanthene-9-carboxylic acid (1.16 g, 0.0051 mol) and dimethylformamide (two drops) in 20 ml of  $\text{CHCl}_3$  (ethanol free) at  $0^\circ\text{C}$ . The mixture was allowed to warm to room temperature under stirring and maintained 1 h at this temperature. After this time, the reaction mixture was concentrated to dryness in vacuo and the obtained residue was dissolved in  $\text{CHCl}_3$  (15 ml) and concentrated again. This procedure was repeated two times. The obtained residue was dissolved in  $\text{CHCl}_3$  and the solution cooled to  $0^\circ\text{C}$ . A solution of 0.865 g (0.0062 mol) of (3R)-N-methylquinuclidin-3-amine (Intermediate I-4) in 10 ml of  $\text{CHCl}_3$  was added. The mixture was allowed to warm to room temperature under stirring. After 4 hours at this temperature, the reaction mixture was treated with  $\text{K}_2\text{CO}_3$  (saturated solution) and the aqueous phase was extracted with  $\text{CHCl}_3$ . The organic solutions were combined and washed with  $\text{K}_2\text{CO}_3$  (saturated solution) and water, dried over  $\text{Na}_2\text{SO}_4$  filtered and concentrated to dryness. The obtained residue was purified by column chromatography (silica gel,  $\text{CHCl}_3$ :  $\text{NH}_4\text{OH}$  99:1  $\rightarrow$   $\text{CHCl}_3$ :  $\text{MeOH}$ :  $\text{NH}_4\text{OH}$  98:2:1 as eluent) to yield 900 mg (50%) of the title product.

MS  $[M+1]^+$ : 349

$^1\text{H}$ -RMN (400 MHz,  $60^\circ\text{C}$ ,  $\text{CDCl}_3$ ):  $\delta$  7.26 - 7.19 (m, 4H), 7.10 - 7.01 (m, 4H), 5.51 (s, 1H), 4.18 (broad multiplet, 1H), 2.92 (s, 3H), 2.85-2.40 (m, 6H), 1.85-1.20 (m, 5H).

**Preparation of (3R)-N-methylquinuclidin-3-amine (Intermediate I-4)**

1g (0.0079 mol) of (3R)-aminoquinuclidine was dissolved in 20 ml of  $\text{CH}_2\text{Cl}_2$  and 1.22 ml (0.0087 mol) of triethylamine were added. The obtained solution was cooled to  $0^\circ\text{C}$ ,  $\text{ClCO}_2\text{Et}$  (0.835 ml, 0.0087 mol) was added and the mixture was stirred 15 h at room temperature. After this time, solvents were evaporated in vacuo, the obtained residue was dissolved in  $\text{CHCl}_3$  and the solution washed with  $\text{K}_2\text{CO}_3$  (saturated solution) and water. The organic phase was dried over  $\text{Na}_2\text{SO}_4$ , filtered and concentrated to dryness. The obtained residue was used without further purification as is described as follows. The obtained residue was dissolved in 15 ml of THF, the solution was cooled to  $0^\circ\text{C}$  and 0.601g (0.016 mol) of  $\text{LiAlH}_4$  were added in several portions. The reaction mixture was heated under reflux for 4h. After this time, was cooled to  $0^\circ\text{C}$  and the excess of hydride was decomposed by the consecutive addition, under stirring, dropwise and cautiously of 0.6 ml of  $\text{H}_2\text{O}$ , 0.6 ml of  $\text{NaOH}$  (10% solution) and 1.8 ml of  $\text{H}_2\text{O}$ . When the

decomposition was finished, the reaction mixture was filtered and the sludge was washed with THF and  $\text{CHCl}_3$ , the organic solutions were combined and concentrated to dryness to obtain 830 mg (75%) of the title product, structure confirmed by  $^1\text{H}$ -RMN and  $^{13}\text{C}$ -RMN.

- 5  $^{13}\text{C}$ -RMN ( $\text{CDCl}_3$ ):  $\delta$  57.0 (CH), 56.5 ( $\text{CH}_2$ ), 47.5 ( $\text{CH}_2$ ), 46.9 ( $\text{CH}_2$ ), 34.3 ( $\text{CH}_3$ ), 26.1 ( $\text{CH}_2$ ), 24.6 (CH), 19.9 ( $\text{CH}_2$ ).

The following Examples of formula (I) were prepared by method (a) from the appropriate carboxylic acid:

- 10 **N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2,3-diphenylpropanamide (Example 25)**  
from 2,3-diphenylpropionic acid.  
**N-(1-Azabicyclo[2.2.2]oct-3-yl)-9H-xanthene-9-carboxamide (Example 28)**  
**N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-9H-xanthene-9-carboxamide (Example 31)**  
**N-[(3S)-1-Azabicyclo[2.2.2]oct-3-yl]-9H-xanthene-9-carboxamide (Example 33)**  
15 All three from 9H-xanthene-9 carboxylic acid.

#### Method (b)

##### **Preparation of N-[(3R)-1-azabicyclo[2.2.2]oct-3-yl]-2,2-dithien-2-ylacetamide**

##### 20 **(Example 17)**

2,2-dithien-2-ylacetic acid was produced by hydrolysis of 2,2-dithien-2-ylacetic acid methyl ester, previously prepared as described in F. Leonard and I. Ehranthal, *J. Am. Chem. Soc.*, (1951), Vol 73, pag 2216. 1.2 g (0.0054 mol) of 2,2-dithien-2-ylacetic acid were dissolved in 25 ml of THF. To this solution were added 0.96 g (0.00594 mol) of 1,1'-

- 25 carbonyldiimidazole and the mixture was refluxed for an hour. The reaction was monitored by TLC following the formation of the imidazolide. When the reaction was completed 0.75 g (0.00594 mol) of (3R)-aminoquinuclidine were added. The reaction mixture was refluxed for 16 h, cooled, diluted with ether and washed with water. The organic layer was extracted with  $\text{HCl}$  2N, the acid solution basified with  $\text{K}_2\text{CO}_3$  and extracted with  $\text{CHCl}_3$ .

- 30 The organic phase was dried over  $\text{Na}_2\text{SO}_4$  and the solvent was evaporated to obtain 0.60 g of an oil which was purified by column chromatography (silica gel,  $\text{CHCl}_3$ :MeOH: $\text{NH}_4\text{OH}$  90:10:1 as eluent). Appropriate fractions were combined and evaporated to give 0.31 g of the title product (17.3%).

mp: 165°C

- 35 MS  $[\text{M}+1]^+$ : 333

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ ) :  $\delta$  1.43 (m, 2H), 1.63 (m, 2H), 1.92 (m, 1H), 2.4 (m, 1H), 2.65-2.85 (m, 4H), 3.31 (m, 1H), 3.97 (m, 1H), 5.37 (s, 1H), 5.98 (d, 1H, NH), 7.0 (m, 4H), 7.28 (m, 2H).

**Preparation of N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-9-hydroxy-9H-fluorene-9-**

**5 carboxamide (Example 44)**

5 g (0.022 mol) of 9-hydroxy-9H-fluorene-9-carboxylic acid were dissolved in 50 ml of dry DMF and 4.2 g (0.026 mol) of 1,1'-carbonyldiimidazole were added. The mixture was stirred during 1 h at room temperature. After this time 3.26 g (0.026 mol) of (3R)-aminoquinuclidine and 0.324 g (0.0027 mol) of DMAP (4-(dimethylamino)pyridine) were  
10 added. The reaction mixture was stirred at room temperature during 14 hours, then was concentrated in vacuo to eliminate the DMF and the obtained residue was dissolved in AcOEt. The organic solution was washed with  $\text{NaHCO}_3$  (saturated solution) and water. The organic layer was separated, dried over  $\text{Na}_2\text{SO}_4$  and solvent was evaporated. The obtained product was purified by column chromatography (silica gel,  $\text{CHCl}_3$ :  $\text{NH}_4\text{OH}$  100:1  
15  $\rightarrow$   $\text{CHCl}_3$ :  $\text{MeOH}$ : $\text{NH}_4\text{OH}$  80:20:1 as eluent). Appropriate fractions were combined and evaporated to yield 304 mg of the title product (4%).

MS  $[\text{M}+1]^+$ : 335

$^1\text{H-RMN}$  ( $\text{CDCl}_3$ ):  $\delta$  7.67 - 7.62 (m, 2H), 7.42 - 7.36 (m, 4H), 7.32 - 7.25 (m, 2H), 6.08 - 5.93 (m, NH, 1H), 3.74 (m, 1H), 3.00 - 2.87 (m, 1H), 2.52 - 2.27 (m, 4H), 2.02 - 1.92 (m,  
20 1H), 1.80 - 1.76 (m, 1H), 1.51 - 1.44 (m, 2H), 1.26 - 1.17 (m, 2H).

**Preparation of (2S)-N-[(3R)-1-azabicyclo[2.2.2]oct-3-yl]-2-cyclopentyl-2-hydroxy-2-thien-2-ylacetamide (Example 49)**

The title compound was prepared as is described in Example 44 starting from (2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-yl acetic acid. The obtained product was purified by  
25 column chromatography (silica gel,  $\text{CHCl}_3$ :  $\text{MeOH}$ :  $\text{NH}_4\text{OH}$  99:1:1  $\rightarrow$   $\text{CHCl}_3$ :  $\text{MeOH}$ : $\text{NH}_4\text{OH}$  97:3:1 as eluent) to give 800 mg of the title product impurified with imidazole. The obtained product was dissolved in  $\text{CHCl}_3$  and washed with  $\text{H}_2\text{O}$  (x 3). The organic solution was dried over  $\text{Na}_2\text{SO}_4$ , filtered and concentrated in vacuo to yield 650  
30 mg of the title product (44% from the starting acid). Structure confirmed by  $^1\text{H-RMN}$  and MS.

MS  $[\text{M}+1]^+$ : 335

$^1\text{H-RMN}$  ( $\text{CDCl}_3$ ):  $\delta$  7.24 (m, 1H), 7.08 (m, 1H), 6.97 (m, 1H), 6.64 (d, NH, 1H), 3.91 - 3.81 (m, 1H), 3.33 - 3.23 (m, 1H), 2.88 - 2.69 (m, 5H), 2.42 - 2.33 (m, 1H), 1.87 (m, 1H),  
35 1.69 - 1.36 (m, 12H)

(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-yl acetic acid has been prepared as is described in WO/02053564

- 5 The following Examples of formula (I) were prepared according to method (b) from the appropriate carboxylic acid:

**N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-cyclopentyl-2-hydroxy-2-thien-2-ylacetamide  
(Example 19)**

- 10 from 2-cyclopentyl-2-hydroxy-2-thien-2-ylacetic acid, which was produced by hydrolysis of 2-cyclopentyl-2-hydroxy-2-thien-2-ylacetic acid methyl ester, compound described in E. Atkinson et al., *J. Med. Chem.*, (1977), Vol 20, n° 12, 1612-1617 and WO02/053564.

**N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-cyclopentyl-2-hydroxy-2-phenylacetamide  
15 (Example 26)**

from 2-cyclopentyl-2-hydroxy-2-phenylacetic acid.

**N-(1-Azabicyclo[2.2.2]oct-3-yl)-9H-xanthene-9-carboxamide (Example 28)**

from 9H-xanthene-9 carboxylic acid

**N-(1-Azabicyclo[2.2.2]oct-3-yl)-9-hydroxy-9H-fluorene-9-carboxamide (Example 41)**

- 20 from 9-Hydroxy-9H-fluorene-9-carboxylic acid.

**Method (c)**

**Preparation of N-[(3R)-1-azabicyclo[2.2.2]oct-3-yl]-2-oxo-2-thien-2-ylacetamide  
25 (Intermediate I-1)**

- Oxalyl chloride (4.5 ml, 0.0516 mol) was added to a solution of 6.68 g (0.0428 mol) of 2-oxo-2-thien-2-ylacetic acid and DMF (several drops) in 100 ml of CHCl<sub>3</sub> (ethanol free) at 0°C. The mixture was stirred and allowed to warm to room temperature. After two hours at this temperature, the solvents were evaporated and the residue was dissolved in CHCl<sub>3</sub> and evaporated again. This procedure was repeated two times. The oil obtained was dissolved in CHCl<sub>3</sub> and the solution obtained cooled at 0°C. A solution of (3R)-aminoquinuclidine (5.91g, 0.0468 mol) in 50 ml of CHCl<sub>3</sub> was added. The mixture was stirred and allowed to warm to room temperature. After 18 hours at this temperature, the reaction mixture was washed with an aqueous K<sub>2</sub>CO<sub>3</sub> solution. The basic aqueous

solution was extracted again with  $\text{CHCl}_3$ . The organic phases were combined, washed with water, dried over  $\text{Na}_2\text{SO}_4$  and evaporated to give 11.34 g of the title product.

MS  $[\text{M}+1]^+$ : 265

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  1.40-1.85 (m, 4H), 2.0 (m, 1H), 2.60 (m, 1H), 2.70-3.0 (m, 4H), 3.4 (m, 1H), 4.0 (m, 1H), 7.20 (m, 1H), 7.50 (d, 1H, NH), 7.85 (m, 1H), 8.40 (m, 1H).

Following an equivalent procedure the following intermediates were prepared:

**N-[(3S)-1-Azabicyclo[2.2.2]oct-3-yl]-2-oxo-2-thien-2-ylacetamide (Intermediate I-2)**

**N-1-Azabicyclo[2.2.2]oct-3-yl]-2-oxo-2-thien-2-ylacetamide (Intermediate I-3)**

10

**Preparation of N-[(3R)-1-azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2,2-dithien-2-yl-acetamide (Example 4)**

A solution of 2-thienylmagnesium bromide was prepared from 2.27 g (0.094 mol) of magnesium and 15.4 g (0.094 mol) of 2-bromothiophene in 150 ml of THF. This solution was added to a solution of 11.34 g (0.043 mol) of N-[(3R)-1-azabicyclo[2.2.2]oct-3-yl]-2-oxo-2-thien-2-ylacetamide (Intermediate I-1) in 120 ml of THF. The mixture was stirred at room temperature for one hour, refluxed for one hour, cooled and treated with a saturated solution of ammonium chloride. The solution obtained was basified with a  $\text{K}_2\text{CO}_3$  solution and extracted with  $\text{CH}_2\text{Cl}_2$ . The organic layer was washed with water and brine, dried over  $\text{Na}_2\text{SO}_4$  and treated with activated carbon. The solution obtained was filtered through Celite and solvents were evaporated to give a solid that was treated with ether and filtered to give 8.77 (58.5%) of the title product.

m. p.: 169°C

MS  $[\text{M}+1]^+$ : 349

$^1\text{H-NMR}$  ( $\text{DMSO}-d_6$ ):  $\delta$  1.30 (m, 1H), 1.42-1.75 (m, 3H), 1.77 (m, 1H), 2.50-2.75 (m, 4H), 2.83 (m, 1H), 3.05 (m, 1H), 3.74 (m, 1H), 6.95 (m, 2H), 7.03 (m, 1H), 7.09 (m, 1H), 7.43 (m, 2H), 7.54 (s, 1H, OH), 7.97 (d, 1H, NH).

The following Examples of formula (I) were prepared according to method (c) using N-[(3R)-1-azabicyclo[2.2.2]oct-3-yl]-2-oxo-2-thien-2-ylacetamide (Intermediate I-1) and the corresponding Grignard reagent:

**N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2-thien-2-ylpent-4-enamide (Example 20).**

**(2\*)-N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2-thien-2-ylbutanamide (diastereomer 1, Example 21) and**

35

**(2\*)-N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2-thien-2-ylbutanamide  
(diastereomer 2, Example 22)**

the two diastereomers were separated by column chromatography (silica gel,  
CHCl<sub>3</sub>/MeOH/NH<sub>4</sub>OH 90:10:1 as eluent). Diastereomer 1: first diastereomer obtained,

5 Diastereomer 2: second distereomer obtained.

**(2\*)-N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2-thien-2-ylbut-3-enamide  
(diastereomer 1, Example 23) and**

**(2\*)-N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2-thien-2-ylbut-3-enamide  
(diastereomer 2, Example 24)**

10 the two diastereomers were separated by column chromatography (silica gel,  
CHCl<sub>3</sub>/MeOH/NH<sub>4</sub>OH 90:10:1 as eluent). Diastereomer 1: first diastereomer obtained,  
Diastereomer 2: second distereomer obtained.

(\*) Configuration not assigned.

15

The following compounds of formula (I) were also prepared according to method (c)

**N-[(3S)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2,2-dithien-2-ylacetamide (Example  
15)**

prepared from Intermediate I-2 and 2-thienyllithium (commercially available).

20 **N-(1-Azabicyclo[2.2.2]oct-3-yl)-2-hydroxy-2,2-dithien-2-ylacetamide (Example 1)**

prepared from Intermediate I-3 and 2-thienyllithium.

**Preparation of N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-(5-bromothien-2-yl)-2-(4-fluoro-  
3-methylphenyl)-2-hydroxyacetamide (Example 48)**

25 Oxalyl chloride (1.16 ml, 0.013 mol) was added to a suspension of 2-(5-bromothien-2-yl)-  
2-oxoacetic acid (2.6 g, 0.011 mol) and two drops of DMF in 30 ml of CHCl<sub>3</sub> (ethanol free)  
cooled to 0°C. The mixture was allowed to warm to room temperature under stirring.

After 1.5 hours the reaction mixture was concentrated to dryness in vacuo and the  
obtained residue was dissolved in CHCl<sub>3</sub> (15 ml) and concentrated again. This procedure

30 was repeated two times. The obtained residue was dissolved in CHCl<sub>3</sub> (30 ml), the  
solution was cooled to 0°C and 1.67 g (0.013 mol) of (3R)-aminoquinuclidine were  
added. The mixture was allowed to warm to room temperature and continued stirring  
during 3 hours. After this time the reaction mixture was treated with K<sub>2</sub>CO<sub>3</sub> (saturated  
solution) and the aqueous layer was extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layers were

35 combined, washed with water, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and evaporated. The obtained



product (N-[(3R)-1-azabicyclo[2.2.2]oct-3-yl]-2-(5-bromothien-2-yl)-2-oxoacetamide) was used without further purification as is described as follows.

The obtained product was dissolved in dry THF (45 ml). The solution obtained was cooled to  $-80^{\circ}\text{C}$  and 14.4 ml of a 1M solution in THF of 4-fluoro-3-methylphenylmagnesium bromide (0.014 mol) were added. The mixture was stirred during 3 hours at low temperature. After this time, the reaction mixture was treated with  $\text{NH}_4\text{Cl}$  (saturated solution) and extracted with  $\text{AcOEt}$ . The organic layer was dried over  $\text{Na}_2\text{SO}_4$ , filtered and solvents were evaporated. The obtained product was purified by column chromatography (silica gel,  $\text{CHCl}_3$ :  $\text{MeOH}$ :  $\text{NH}_4\text{OH}$  99:1:1  $\rightarrow$   $\text{CHCl}_3$ :  $\text{MeOH}$ :  $\text{NH}_4\text{OH}$  96:4:1) to yield 2.1g of the title compound (42% from the starting acid) as a mixture of diastereomers.

MS  $[\text{M}+1]^+$ : 453, 455

$^1\text{H}$ -RMN ( $\text{CDCl}_3$ ): (mixture of diastereomers)  $\delta$  7.35 - 7.20 (m, 2H), 7.08 - 6.89 (m, 3H), 6.80 (dd, 1H), 3.93 - 3.84 (m, 1H), 3.18 - 3.07 (m, 1H), 2.69 - 2.58 (m, 4H), 2.33 - 2.22 (m, 4H), 1.92 - 1.87 (m, 1H), 1.63 - 1.42 (m, 4H)

2-(5-bromothien-2-yl)-2-oxoacetic acid was obtained by hydrolysis of ethyl 2-(5-bromothien-2-yl)-2-oxoacetate by a standard method ( $\text{EtOH}$ ,  $\text{NaOH}$  2N,  $60^{\circ}\text{C}$ , 1.5 h).

#### Preparation of N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-fur-2-yl-2-hydroxypent-3-ynamide (Example 45)

To a suspension cooled to  $0^{\circ}\text{C}$  of 1.72 g (0.012 mol) of 2-fur-2-yl-2-oxoacetic acid and two drops of DMF in 40 ml of  $\text{CHCl}_3$  (ethanol free), 1.17 ml (0.013 mol) of oxalyl chloride were added. The resulting mixture was allowed to warm to room temperature under stirring and continued stirring during 1 hour. After this time, the mixture was concentrated to dryness in vacuo and the obtained residue was dissolved in  $\text{CHCl}_3$  (20 ml) and concentrated again. This procedure was repeated two times. The obtained product was dissolved in  $\text{CHCl}_3$  (50 ml), the solution was cooled to  $0^{\circ}\text{C}$  and 1.7 g (0.013 mol) of (3R)-aminoquinuclidine were added. The mixture was allowed to warm to room temperature and continued stirring during 16 hours. After this time the reaction mixture was treated with  $\text{K}_2\text{CO}_3$  (saturated solution) and the aqueous layer was extracted with  $\text{CH}_2\text{Cl}_2$ . The organic layers were combined, washed with water, dried over  $\text{Na}_2\text{SO}_4$ , filtered and evaporated. The obtained product (N-[(3R)-1-azabicyclo[2.2.2]oct-3-yl]-2-(2-furyl)-2-oxoacetamide) was used without further purification as is described as follows.

This product was dissolved in dry THF (50 ml). The solution obtained was cooled to  $-80^{\circ}\text{C}$  and 28 ml of a 0.5 M solution in THF of 1-propynylmagnesium bromide (0.014 mol) were

added. The mixture was allowed to warm to room temperature during 3 hours. After this time, the reaction mixture was treated with  $\text{NH}_4\text{Cl}$  (saturated solution) and extracted with  $\text{AcOEt}$  and  $\text{CH}_2\text{Cl}_2$ . The organic layers were combined, dried over  $\text{Na}_2\text{SO}_4$ , filtered and solvents were evaporated. The obtained product was purified by column chromatography (silica gel,  $\text{CHCl}_3$ :  $\text{MeOH}$ :  $\text{NH}_4\text{OH}$  97:3:1  $\rightarrow$   $\text{CHCl}_3$ :  $\text{MeOH}$ :  $\text{NH}_4\text{OH}$  94:6:1 as eluent) to yield 1.38 g of the title compound (39% from the starting acid) as a mixture of diastereomers.

MS  $[\text{M}+1]^+$ : 289

$^1\text{H}$ -RMN ( $\text{CDCl}_3$ ): (mixture of diastereomers)  $\delta$  7.36 - 7.35 (m, 1H), 6.87 - 6.76 (m, NH, 1H), 6.54 - 6.52 (m, 1H), 6.34 - 6.32 (m, 1H), 3.92 (m, 1H), 3.29 - 3.17 (m, 1H), 2.78 - 2.64 (m, 4H), 2.47 - 2.35 (m, 1H), 2.00 - 1.90 (m, 1H), 1.92 (s, 3H), 1.70 - 1.39 (m, 4H).

**Preparation of N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-fur-2-yl-2-hydroxy-4-(4-methoxyphenyl)butanamide (Example 46)**

This product was prepared according to the procedure described in Example 45, starting from 2 g (0.014 mol) of 2-fur-2-yl-2-oxoacetic acid, and using 4-methoxyphenethylmagnesium chloride (0.021 mol, 43 ml of a 0.5 M solution in THF) as Grignard reagent. The obtained product was purified by column chromatography (silica gel,  $\text{CHCl}_3$ :  $\text{MeOH}$ :  $\text{NH}_4\text{OH}$  99:1:1  $\rightarrow$   $\text{CHCl}_3$ :  $\text{MeOH}$ :  $\text{NH}_4\text{OH}$  97:3:1 as eluent) to yield 2.4 g of the title compound (44% from the starting acid) as a mixture of diastereomers.

MS  $[\text{M}+1]^+$ : 385

$^1\text{H}$ -RMN ( $\text{CDCl}_3$ ): (mixture of diastereomers)  $\delta$  7.39 - 7.38 (m, 1H), 7.09 (d, 2H), 6.82 (d, 2H), 6.70 - 6.67 (m, NH, 1H), 6.41 - 6.35 (m, 2H), 3.92 (m, 1H), 3.78 (s, 3H), 3.33 - 3.24 (m, 1H), 2.82 - 2.19 (m, 9H), 1.96 - 1.87 (m, 1H), 1.67 - 1.40 (m, 4H)

25

**Method (d)**

**Preparation of (3R)-1-(3-phenoxypropyl)-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane bromide (Example 32)**

0.3 g (0.00089 mol) of N-[(3R)-1-azabicyclo[2.2.2]oct-3-yl]-9H-xanthene-9-carboxamide (Example 31) were dissolved in 6 ml of THF. To this solution 0.21 ml (0.287g, 0.001335 mol) of (3-bromopropoxy)benzene were added. The mixture was refluxed during 4 hours and then stirred 17 h at room temperature. The reaction mixture was filtered and the solid obtained washed with ether several times and dried. 0.48 g (98%) of the title product were obtained.

35

m.p.: 278.5-279.2°C

MS [M-Br]<sup>+</sup>: 469

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 1.80-2.0 (m, 3H), 2.05-2.30 (m, 4H), 3.15-3.25 (m, 1H), 3.30-3.65 (m, 6H), 3.82 (m, 1H), 4.05 (t, 2H), 4.10 (m, 1H), 5.05 (s, 1H), 6.90-7.0 (m, 3H), 7.05-7.20 (m, 4H), 7.25-7.42 (m, 6H), 9.01 (d, 1H, NH).

**Preparation of 3-[[[9-hydroxy-9H-fluoren-9-yl]carbonyl]amino]-1-methyl-1-azoniabicyclo[2.2.2]octane bromide (Example 42)**

0.245 g (0.00073 mol) of N-(1-azabicyclo[2.2.2]oct-3-yl)-9-hydroxy-9H-fluorene-9-carboxamide (Example 41) were dissolved in 4 ml of acetonitrile and 6 ml of CHCl<sub>3</sub>. To this solution 5.46 ml of a 1 M solution of methyl bromide in acetonitrile were added and the mixture was stirred for 96 h at room temperature under N<sub>2</sub> atmosphere. After this time the solvents were evaporated. The residue was coevaporated with ether three times. Ether was added to the residue and the mixture stirred. The solid obtained was filtered and washed several times with ether. The yield was 0.26 g (83.8%) of the title compound. m.p.: 197.5-203.6°C

MS [M-Br]<sup>+</sup>: 349

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 1.80-2.0 (m, 3H), 2.10-2.25 (m, 2H), 2.96 (s, 3H), 3.35-3.70 (m, 5H), 3.78 (m, 1H), 4.15 (m, 1H), 6.82 (s, 1H, OH), 7.30 (m, 2H), 7.43 (m, 4H), 7.79 (m, 2H), 8.81 (d, 1H, NH).

**Method (e)**

**Preparation of (3S)-1-cyclohexylmethyl-3-[[9H-xanthen-9-yl]carbonyl]amino]-1-azoniabicyclo[2.2.2]octane trifluoroacetate (Example 36)**

28.15 mg (0.0842 mmol) of N-[(3S)-1-azabicyclo[2.2.2]oct-3-yl]-9H-xanthene-9-carboxamide (Example 33) were dissolved in 0.5 ml of DMSO. A solution of 74.37 mg (0.421 mmol) of (bromomethyl)cyclohexane in 0.5 ml of DMSO was added and the mixture stirred at room temperature overnight. The mixture was purified by solid phase extraction with a cation exchange Mega Bond Elut cartridge, previously conditioned at pH=7.5 with 0.1 M NaH<sub>2</sub>PO<sub>4</sub> buffer. The reaction mixture was applied to the cartridge and washed first with 2 ml of DMSO and then three times with 5 ml of acetonitrile, rinsing away all starting materials. The ammonium derivative was eluted with 5 ml of 0.03 M TFA solution in CH<sub>3</sub>CN:CHCl<sub>3</sub> (2:1). This solution was neutralized with 300 mg of poly(4-

vinylpyridine), filtered and evaporated to dryness. The yield was 12.4 mg of the title product. (27.0%).

[M-CF<sub>3</sub>COO]<sup>+</sup>: 431

**5 Preparation of (3R)-3-(2-hydroxy-2,2-dithien-2-ylacetyl-amino)-1-[3-(3-hydroxyphenoxy)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate (Example 12)**

The title compound was prepared as described for Example 36 from 30 mg (0.0861 mmol) of N-[(3R)-1-azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2,2-dithien-2-ylacetamide (Example 4) (dissolved in 0.5 ml of DMSO) and 78 mg (0.418 mmol) of 3-(3-chloropropoxy)phenol (dissolved in 0.5 ml of DMSO). The yield was 12.7 mg of the title product. (24%).

[M-CF<sub>3</sub>COO]<sup>+</sup>: 499

Also included within the scope of the present invention are pharmaceutical compositions which comprise, as the active ingredient, at least one quinuclidine amide derivative of formulae (I) or (II) in association with a pharmaceutically acceptable carrier or diluent. Preferably the composition is made up in a form suitable for oral administration.

The pharmaceutically acceptable carrier or diluents which are mixed with the active compound or compounds, to form the composition of this invention are well-known per se and the actual excipients used depend inter alia on the intended method of administration of the composition.

Compositions of this invention are preferably adapted for oral administration. In this case, the composition for oral administration may take the form of tablets, film-coated tablets, liquid inhalant, powder inhalant and inhalation aerosol; all containing one or more compounds of the invention; such preparations may be made by methods well-known in the art.

The diluents which may be used in the preparations of the compositions include those liquid and solid diluents which are compatible with the active ingredient, together with colouring or flavouring agents, if desired. Tablets or film-coated tablets may conveniently contain between 1 and 500 mg, preferably from 5 to 300 mg of active ingredient. The inhalant compositions may contain between 1 µg and 1,000 µg, preferably from 10 µg to 800 µg of active ingredient. In human therapy, the dose of the compound of formulae (I) or

(II) depend on the desired effect and duration of treatment; adult doses are generally between 3 mg and 300 mg per day as tablets and 10 µg and 800 µg per day as inhalant composition.

## 5 Pharmacological Action

The results on human muscarinic receptors binding and in the test on bronchospasm in guinea pig, were obtained as described below.

### 10 **Human muscarinic receptor studies.**

The binding of [3H]-NMS to human muscarinic receptors was performed according to Waelbroeck et al (1990), Mol. Pharmacol., 38: 267-273. Assays were carried out at 25°C. Membrane preparations from stably transfected chinese hamster ovary-K1 cells (CHO) expressing the genes for the human M3 muscarinic receptors were used.

For determination of IC<sub>50</sub>, membrane preparations were suspended in DPBS to a final concentration of 89 µg/ml for the M3 subtype. The membrane suspension was incubated with the tritiated compound for 60 min. After incubation the membrane fraction was separated by filtration and the bound radioactivity determined. Non-specific binding was determined by addition of 10<sup>-4</sup> M atropine. At least six concentrations were assayed in duplicate to generate individual displacement curves.

Our results show that the compounds of the present invention have high affinities for M3 muscarinic receptors, preferably human muscarinic receptors. Thus, the IC<sub>50</sub> of the preferred compounds of the invention is lower than 100 nM. Most preferred compounds, such as the compounds of examples 1, 3, 4, 8, 10, 11, 14, 19, 30 and 32 described below, have an IC<sub>50</sub> lower than 60 nM.

### 30 **Test on bronchospasm in guinea pig**

The studies were performed according to H. Konzett and F. Rössler (1940), Arch. Exp. Path. Pharmacol. 195: 71-74. Aqueous solutions of the agents to be tested were nebulized and inhaled by anaesthetized ventilated male guinea pigs (Dunkin-Hartley). Bronchial response to intravenous acetylcholine challenge was determined before and

after drug administration and the changes in pulmonary resistance at several time-points were expressed as percent of inhibition of bronchospasm.

The compounds of the present invention inhibited the bronchospasm response to acetylcholine with high potency and a long duration of action.

From the above described results one of ordinary skill in the art can readily understand that the compounds of the present invention have excellent M3 antimuscarinic activity and thus are useful for the treatment of diseases in which the M3 muscarinic receptor is implicated, including respiratory disorders such as chronic obstructive pulmonary disease (COPD), bronchitis, bronchial hyperreactivity, asthma, cough and rhinitis; urological disorders such as urinary incontinence, pollakiuria, neurogenic or unstable bladder, cystospasm and chronic cystitis; gastrointestinal disorders such as irritable bowel syndrome, spastic colitis, diverticulitis and peptic ulceration; and cardiovascular disorders such as vagally induced sinus bradycardia.

The present invention further provides a compound of formulae (I) or (II) or a pharmaceutically acceptable composition comprising a compound of formulae (I) or (II) for use in a method of treatment of the human or animal body by therapy, in particular for the treatment of respiratory, urological or gastrointestinal disease or disorder.

The present invention further provides the use of a compound of formulae (I) or (II) or a pharmaceutically acceptable composition comprising a compound of formulae (I) or (II) for the manufacture of a medicament for the treatment of a respiratory, urological or gastrointestinal disease or disorder.

Further, the compounds of formulae (I) or (II) and pharmaceutical compositions comprising a compound of formulae (I) or (II) can be used in a method of treating a respiratory, urological or gastrointestinal disease or disorder, which method comprises administering to a human or animal patient in need of such treatment an effective, non-toxic, amount of a compound of formulae (I) or (II) or a pharmaceutical composition comprising a compound of formulae (I) or (II).

Further, the compounds of formulae (I) or (II) and pharmaceutical compositions comprising a compound of formulae (I) or (II) can be used in combination with other drugs

effective in the treatment of these diseases. For example with  $\beta_2$  agonists, steroids, antiallergic drugs; phosphodiesterase IV inhibitors and/or leukotriene D4 (LTD4) inhibitors, for simultaneous, separate or sequential use in the treatment of a respiratory disease.

- 5 The present invention will be further illustrated by the following examples. The examples are given by way of illustration only and are not to be construed as a limiting.

#### Example 1

##### **N-(1-Azabicyclo[2.2.2]oct-3-yl)-2-hydroxy-2,2-dithien-2-ylacetamide**

- 10 The title compound was synthesised according to method c. The yield was 0.67 g, 15.2%.  
m.p.: 185°C.

MS [M+1]<sup>+</sup>: 349.

- <sup>1</sup>H-NMR (DMSO-d<sub>6</sub>):  $\delta$  1.30 (m, 1H), 1.40-1.75 (m, 3H), 1.77 (m, 1H), 2.50-2.75 (m, 4H),  
2.83 (m, 1H), 3.05 (m, 1H), 3.74 (m, 1H), 6.94 (m, 2H), 7.04 (m, 1H), 7.10 (m, 1H), 7.43  
15 (m, 2H), 7.53 (s, 1H, OH), 7.95 (d, 1H, NH).

#### Example 2

##### **3-(2-Hydroxy-2,2-dithien-2-ylacetyl-amino)-1-methyl-1-azoniabicyclo[2.2.2]octane bromide**

- 20 The title compound was synthesised according to methods c and d. The yield of the final step was 0.16 g, 68%.

MS [M-Br]<sup>+</sup>: 363.

- <sup>1</sup>H-NMR (DMSO-d<sub>6</sub>):  $\delta$  1.76 (m, 1H), 1.85-2.08 (m, 3H), 2.14 (m, 1H), 2.92 (s, 3H), 3.30-  
3.60 (m, 5H), 3.75 (m, 1H), 4.20 (s, 1H), 7.0 (m, 2H), 7.06 (m, 1H), 7.12 (m, 1H), 7.46 (m,  
25 2H), 7.65 (s, 1H, OH), 8.64 (d, 1H, NH).

#### Example 3

##### **3-(2-Hydroxy-2,2-dithien-2-ylacetyl-amino)-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane bromide**

- 30 The title compound was synthesised according to methods c and d. The yield of the final step was 0.24 g, 82.7%.

m.p.: 180.6-188.3°C.

MS [M-Br]<sup>+</sup>: 483.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 1.78 (m, 1H), 1.90-2.25 (m, 6H), 3.30-3.65 (m, 7H), 3.67-3.80 (m, 1H), 4.05 (t, 2H), 4.24 (m, 1H), 6.94-7.0 (m, 5H), 7.07 (m, 1H), 7.13 (m, 1H), 7.31 (m, 2H), 7.47 (m, 2H), 7.66 (s, 1H, OH), 8.74 (d, 1H, NH).

5 **Example 4**

**N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2,2-dithien-2-ylacetamide**

The title compound was synthesised according to method c. The yield was 8.77 g, 58.5%.  
m.p.: 169°C.

MS [M+1]<sup>+</sup>: 349.

10 <sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 1.30 (m, 1H), 1.42-1.75 (m, 3H), 1.77 (m, 1H), 2.50-2.75 (m, 4H), 2.83 (m, 1H), 3.05 (m, 1H), 3.74 (m, 1H), 6.95 (m, 2H), 7.03 (m, 1H), 7.09 (m, 1H), 7.43 (m, 2H), 7.54 (s, 1H, OH), 7.97 (d, 1H, NH).

**Example 5**

15 **(3R)-3-(2-Hydroxy-2,2-dithien-2-ylacetyl-amino)-1-methyl-1-azoniabicyclo[2.2.2]octane trifluoroacetate**

The title compound was synthesised according to methods c and e. The yield of the final step was 0.0124 g, 30.2%.

MS [M-CF<sub>3</sub>COO]<sup>+</sup>: 363.

20

**Example 6**

**(3R)-1-Allyl-3-(2-hydroxy-2,2-dithien-2-ylacetyl-amino)-1-azoniabicyclo[2.2.2]octane trifluoroacetate**

25 The title compound was synthesised according to methods c and e. The yield of the final step was 0.010 g, 23.1%.

MS [M-CF<sub>3</sub>COO]<sup>+</sup>: 389.

**Example 7**

30 **(3R)-1-Heptyl-3-(2-hydroxy-2,2-dithien-2-ylacetyl-amino)-1-azoniabicyclo[2.2.2]octane trifluoroacetate**

The title compound was synthesised according to methods c and e. The yield of the final step was 0.0118 g, 24.4%.

MS [M-CF<sub>3</sub>COO]<sup>+</sup>: 447.



**Example 8****(3R)-3-(2-Hydroxy-2,2-dithien-2-ylacetyl-amino)-1-(3-phenylpropyl)-1-azoniabicyclo[2.2.2]octane bromide**

The title compound was synthesised according to methods c and d. The yield of the final  
5 step was 0.49 g, 78%.

m.p.: 117.3-118.9°C.

MS [M-Br]<sup>+</sup>: 467

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 1.75 (m, 1H), 1.80-2.05 (m, 5H), 2.09 (m, 1H), 2.60 (m, 2H), 3.16  
(m, 2H), 3.25-3.60 (m, 5H), 3.68 (m, 1H), 4.21 (m, 1H), 6.96 (m, 2H), 7.04 (m, 1H), 7.11  
10 (m, 1H), 7.20-7.40 (m, 5H), 7.45 (m, 2H), 7.62 (s, 1H, OH), 8.70 (d, 1H, NH).

**Example 9****(3R)-3-(2-Hydroxy-2,2-dithien-2-ylacetyl-amino)-1-((E)-3-phenylallyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate**

15 The title compound was synthesised according to methods c and e. The yield of the final step was 0.0032 g, 6.4%.

[M-CF<sub>3</sub>COO]<sup>+</sup>: 465.

**Example 10****(3R)-3-(2-Hydroxy-2,2-dithien-2-ylacetyl-amino)-1-(2-phenoxyethyl)-1-azoniabicyclo[2.2.2]octane bromide**

The title compound was synthesised according to methods c and d. The yield of the final  
step was 0.470 g, 74%.

m.p.: 112.6-113.9°C.

25 MS [M-Br]<sup>+</sup>: 469

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 1.77 (m, 1H), 1.90-2.15 (m, 4H), 3.40-3.80 (m, 7H), 3.86 (m, 1H),  
4.24 (m, 1H), 4.43 (m, 2H), 6.95-7.0 (m, 5H), 7.04 (m, 1H), 7.11 (m, 1H), 7.35 (m, 2H),  
7.46 (m, 2H), 7.66 (s, 1H, OH), 8.72 (d, 1H, NH).

**30 Example 11****(3R)-3-(2-Hydroxy-2,2-dithien-2-ylacetyl-amino)-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane bromide**

The title compound was synthesised according to methods c and d. The yield of the final  
step was 0.65 g, 80%.

35 m.p.: 182°C.

MS [M-Br]<sup>+</sup>: 483

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 1.76 (m, 1H), 1.85-2.25 (m, 6H), 3.25-3.65 (m, 7H), 3.65-3.82 (m, 1H), 4.05 (t, 2H), 4.23 (m, 1H), 6.90-7.0 (m, 5H), 7.04 (m, 1H), 7.11 (m, 1H), 7.31 (m, 2H), 7.45 (m, 2H), 7.65 (s, 1H, OH), 8.73 (d, 1H, NH).

5

#### Example 12

**(3R)-3-(2-Hydroxy-2,2-dithien-2-ylacetyl-amino)-1-[3-(3-hydroxyphenoxy)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate**

The title compound was synthesised according to methods c and e. The yield of the final step was 0.0127 g, 24.0%.

[M-CF<sub>3</sub>COO]<sup>+</sup>: 499.

#### Example 13

**(3R)-1-(2-Benzyloxyethyl)-3-(2-hydroxy-2,2-dithien-2-ylacetyl-amino)-1-azoniabicyclo[2.2.2]octane trifluoroacetate**

The title compound was synthesised according to methods c and e. The yield of the final step was 0.0146 g, 28.4%.

[M-CF<sub>3</sub>COO]<sup>+</sup>: 483.

#### Example 14

**(3R)-3-(2-Hydroxy-2,2-dithien-2-ylacetyl-amino)-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane bromide**

The title compound was synthesised according to methods c and d. The yield of the final step was 0.49 g, 77%.

m.p.: 111.1-113.2°C.

MS [M-Br]<sup>+</sup>: 473

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 1.75 (m, 1H), 1.85-2.15 (m, 6H), 2.84 (t, 2H), 3.17 (m, 2H), 3.25-3.45 (m, 3H), 3.45-3.62 (m, 2H), 3.70 (m, 1H), 4.21 (m, 1H), 6.90-7.0 (m, 4H), 7.05 (m, 1H), 7.11 (m, 1H), 7.38 (m, 1H), 7.45 (m, 2H), 7.65 (s, 1H, OH), 8.71 (d, 1H, NH).

30

#### Example 15

**N-[(3S)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2,2-dithien-2-ylacetamide**

The title compound was synthesised according to method c. The yield was 0.44 g, 7.67%.

MS [M+1]<sup>+</sup>: 349.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.40-1.70 (m, 4H), 1.98 (m, 1H), 2.40-2.47 (m, 1H), 2.60-2.85 (m, 4H), 3.24 (m, 1H), 3.97 (m, 1H), 6.91 (d, 1H), 7.01 (m, 2H), 7.15 (m, 2H), 7.33 (m, 2H).

#### Example 16

5 **(3S)-3-(2-Hydroxy-2,2-dithien-2-ylacetyl-amino)-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane bromide**

The title compound was synthesised according to methods c and d. The yield of the final step was 0.36 g, 70.6%.

m.p.: 172.8-173.9°C.

10 MS [M-Br]<sup>+</sup>: 483.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 1.78 (m, 1H), 1.85-2.25 (m, 6H), 3.25-3.65 (m, 7H), 3.70-3.82 (m, 1H), 4.04 (t, 2H), 4.24 (m, 1H), 6.90-7.0 (m, 5H), 7.06 (m, 1H), 7.12 (m, 1H), 7.31 (m, 2H), 7.45 (m, 2H), 7.65 (s, 1H, OH), 8.74 (d, 1H, NH).

#### 15 Example 17

**N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2,2-dithien-2-ylacetamide**

The title compound was synthesised according to method b. The yield was 0.31 g, 17.3%.

m.p.: 165°C.

MS [M+1]<sup>+</sup>: 333.

20 <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.43 (m, 2H), 1.63 (m, 2H), 1.92 (m, 1H), 2.4 (m, 1H), 2.65-2.85 (m, 4H), 3.31 (m, 1H), 3.97 (m, 1H), 5.37 (s, 1H), 5.98 (d, 1H, NH), 7.0 (m, 4H), 7.28 (m, 2H).

#### Example 18

25 **(3R)-3-(2,2-Dithien-2-ylacetyl-amino)-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane bromide**

The title compound was synthesised according to methods b and d. The yield of the final step was 0.21 g, 79.8%.

m.p.: 135.6-137.1°C.

MS [M-Br]<sup>+</sup>: 467.

30 <sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 1.83 (m, 1H), 1.94 (m, 2H), 2.0-2.25 (m, 4H), 3.19 (m, 1H), 3.30-3.55 (m, 6H), 3.86 (m, 1H), 4.03 (t, 2H), 4.18 (m, 1H), 5.65 (s, 1H), 6.93-6.98 (m, 5H), 7.01-7.06 (m, 2H), 7.31 (m, 2H), 7.42 (m, 2H), 9.10 (d, 1H, NH).

**Example 19****N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-cyclopentyl-2-hydroxy-2-thien-2-ylacetamide**

The title compound was synthesised as a mixture of diastereomers according to method

b. The yield was 0.12 g, 27.9%.

5 MS [M+1]<sup>+</sup>: 335.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): (mixture of diastereomers 50:50) δ 1.20-1.80 (m, 13H), 2.50-2.90 (m, 6H), 3.04 (m, 1H), 3.69 (m, 1H), 5.98 & 6.01 (s, 1H, OH), 6.93 (m, 1H), 7.08 (m, 1H), 7.35 (m, 1H), 7.59 (m, 1H, NH).

10 **Example 20**

**N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2-thien-2-ylpent-4-enamide**

The title compound was synthesised as a mixture of diastereomers according to method

c. The yield was 1.92 g, 82.4%.

m.p.: 54.4-58.3°C

15 MS [M+1]<sup>+</sup>: 307.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): (mixture of diastereomers) δ 1.25 (m, 1H), 1.40-1.75 (m, 4H), 2.45-2.70 (m, 5H), 2.70-3.10 (m, 3H), 3.65 (m, 1H), 5.0-5.10 (m, 2H), 5.60-5.80 (m, 1H), 6.50 & 6.52 (s, 1H, OH), 6.95 (m, 1H), 7.05 (m, 1H), 7.38 (m, 1H), 7.60 (d, 1H, NH).

20 **Example 21**

**(2\*)-N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2-thien-2-ylbutanamide**

(diastereomer 1, 70:30)

The title compound was synthesised according to method c. The yield was 0.19 g, 34% (based on single isomer).

25 m.p.: 139.0-140.7°C.

MS [M+1]<sup>+</sup>: 295.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): (diastereomer 1, 70:30) δ 0.80 (t, 3H), 1.27 (m, 1H), 1.40-1.80 (m, 4H), 1.83 (m, 1H), 2.16 (m, 1H), 2.45-2.70 (m, 4H), 2.79 (m, 1H), 2.90-3.10 (m, 1H), 3.67 (m, 1H), 6.28 & 6.31 (s, 1H, OH), 6.94 (m, 1H), 7.03 (m, 1H), 7.36 (m, 1H), 7.62 (d, 1H,

30 NH).

(\*): Configuration not assigned.

**Example 22****(2\*)-N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2-thien-2-ylbutanamide**

35 (diastereomer 2, 27:73)

The title compound was synthesised according to method c. The yield was 0.42 g, 75% (based on single isomer).

m.p.: 68.9-70.2°C.

MS [M+1]<sup>+</sup>: 295.

5 <sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): (diastereomer 2, 27:73) δ 0.82 (t, 3H), 1.26 (m, 1H), 1.40-1.80 (m, 4H), 1.86 (m, 1H), 2.17 (m, 1H), 2.45-2.70 (m, 4H), 2.78 (m, 1H), 2.90-3.10 (m, 1H), 3.67 (m, 1H), 6.29 & 6.32 (s, 1H, OH), 6.93 (m, 1H), 7.03 (m, 1H), 7.36 (m, 1H), 7.62 (d, 1H, NH).

(\*): Configuration not assigned.

10

#### Example 23

**(2\*)-N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2-thien-2-ylbut-3-enamide**

(diastereomer 1)

The title compound was synthesised according to method c. The yield was 0.21 g, 18.9 %

15 (based on single isomer).

m.p.: 171.7-173.2°C.

MS [M+1]<sup>+</sup>: 293.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): (diastereomer 1) δ 1.27 (m, 1H), 1.40-1.75 (m, 4H), 2.50-2.70 (m, 4H), 2.77 (m, 1H), 3.0 (m, 1H), 3.66 (m, 1H), 5.18 (d, 1H), 5.38 (d, 1H), 6.41 (dd, 1H),  
20 6.71 (s, 1H, OH), 6.95 (m, 1H), 7.02 (m, 1H), 7.41 (m, 1H), 7.70 (d, 1H, NH).

(\*): Configuration not assigned.

#### Example 24

**(2\*)-N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2-thien-2-ylbut-3-enamide**

25 (diastereomer 2)

The title compound was synthesised according to method c. The yield was 0.21 g, 18.9 %

(based on single isomer)

m.p.: 53.8-55.0°C.

MS [M+1]<sup>+</sup>: 293.

30 <sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): (diastereomer 2) δ 1.27 (m, 1H), 1.40-1.75 (m, 4H), 2.50-2.70 (m, 4H), 2.81 (m, 1H), 3.03 (m, 1H), 3.69 (m, 1H), 5.21 (d, 1H), 5.40 (d, 1H), 6.45 (dd, 1H), 6.74 (s, 1H, OH), 6.96 (m, 1H), 7.01 (m, 1H), 7.41 (m, 1H), 7.72 (d, 1H, NH).

(\*): Configuration not assigned.

**Example 25****N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2,3-diphenylpropanamide**

The title compound was synthesised as a mixture of diastereomers according to method

a. The yield was 1.21 g, 82.3 %.

5 m.p.: 142°C.

MS [M+1]<sup>+</sup>: 335.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): (mixture of diastereomers) δ 0.95-1.30 (m, 2H), 1.35-1.50 (m, 2H), 1.52 (m, 1H), 2.14-2.27 (m, 1H), 2.50-2.65 (m, 4H), 2.81-2.96 (m, 2H), 3.24-3.32 (m, 1H), 3.56 (m, 1H), 3.78-3.89 (m, 1H), 7.10-7.45 (m, 10H), 8.13 (m, 1H, NH).

10

**Example 26****N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-cyclopentyl-2-hydroxy-2-phenylacetamide**

The title compound was synthesised as a mixture of diastereomers according to method

b. The yield was 0.25 g, 28.4 %.

15 m.p.: 69.8-73.3°C.

MS [M+1]<sup>+</sup>: 329.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): (mixture of diastereomers) δ 1.25 (m, 3H), 1.35-1.75 (m, 10H), 2.40-2.70 (m, 4H), 2.76 (m, 1H), 2.90-3.10 (m, 2H), 3.62 (m, 1H), 5.60 & 5.62 (s, 1H, OH), 7.20-7.23 (m, 1H), 7.27-7.33 (m, 2H), 7.52-7.61 (m, 3H).

20

**Example 27****N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-phenylhexanamide**

The title compound was synthesised as a mixture of diastereomers according to method

a. The yield was 1.59 g, 63.1 %.

25 m.p.: 136°C.

MS [M+1]<sup>+</sup>: 301.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): (mixture of diastereomers 50:50) δ 0.84 (t, 3H), 1.10-1.60 (m, 9H), 1.70 (m, 1H), 1.90 (m, 1H), 2.25-2.50 (m, 1H), 2.50-2.80 (m, 4H), 2.94 & 3.05 (m, 1H), 3.50 (m, 1H), 3.65 (m, 1H), 7.15-7.40 (m, 5H), 8.04 (m, 1H, NH).

30

**Example 28****N-(1-Azabicyclo[2.2.2]oct-3-yl)-9H-xanthene-9-carboxamide**

The title compound was synthesised according to method b. The yield was 0.28g, 19 %.

m.p.: 251°C.

35 MS [M+1]<sup>+</sup>: 335.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 1.25-1.60 (m, 3H), 1.67 (m, 1H), 1.82 (m, 1H), 2.40-2.50 (m, 1H), 2.55-2.75 (m, 3H), 2.82 (m, 1H), 3.06 (m, 1H), 3.62 (m, 1H), 5.0 (s, 1H), 7.05-7.15 (m, 4H), 7.25-7.35 (m, 4H), 8.51 (d, 1H, NH).

The title compound was also synthesised according to method a. The yield was 2.11 g, 52.6 %.

#### Example 29

##### 1-Methyl-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane bromide

The title compound was synthesised according to methods a and d. The yield of the final step was 0.25 g, 77.6%.

m.p.: 318°C.

MS [M-Br]<sup>+</sup>: 349.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 1.80-2.0 (m, 3H), 2.10 (m, 1H), 2.22 (m, 1H), 2.95 (s, 3H), 3.10-3.20 (m, 1H), 3.30-3.50 (m, 4H), 3.80 (m, 1H), 4.05 (m, 1H), 5.06 (s, 1H), 7.08-7.16 (m, 4H), 7.28-7.35 (m, 4H), 8.98 (d, 1H, NH).

#### Example 30

##### 1-(3-Phenoxypropyl)-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane bromide

The title compound was synthesised according to methods a and d. The yield of the final step was 0.44 g, 100%.

m.p.: 242°C.

MS [M-Br]<sup>+</sup>: 469.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 1.80-2.0 (m, 3H), 2.05-2.30 (m, 4H), 3.25-3.70 (m, 7H), 3.82 (m, 1H), 4.05 (t, 2H), 4.10 (m, 1H), 5.15 (s, 1H), 6.90-7.0 (m, 3H), 7.05-7.20 (m, 4H), 7.25-7.42 (m, 6H), 9.16 (d, 1H, NH).

#### Example 31

##### N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-9H-xanthene-9-carboxamide

The title compound was synthesised according to method a. The yield was 1.03 g, 43.4 %.

m.p.: 239.7-241.4°C.

MS [M+1]<sup>+</sup>: 335.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 1.25-1.60 (m, 3H), 1.66 (m, 1H), 1.82 (m, 1H), 2.42-2.50 (m, 1H), 2.55-2.75 (m, 3H), 2.85 (m, 1H), 3.05 (m, 1H), 3.60 (m, 1H), 5.0 (s, 1H), 7.05-7.15 (m, 4H), 7.25-7.35 (m, 4H), 8.52 (d, 1H, NH).

5

**Example 32**

**(3R)-1-(3-Phenoxypropyl)-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane bromide**

The title compound was synthesised according to methods a and d. The yield of the final  
10 step was 0.48 g, 98%.

m.p.: 278.5-279.2°C.

MS [M-Br]<sup>+</sup>: 469.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 1.80-2.0 (m, 3H), 2.05-2.30 (m, 4H), 3.15-3.25 (m, 1H), 3.30-3.65  
(m, 6H), 3.82 (m, 1H), 4.05 (t, 2H), 4.10 (m, 1H), 5.05 (s, 1H), 6.90-7.0 (m, 3H), 7.05-7.20  
15 (m, 4H), 7.25-7.42 (m, 6H), 9.01 (d, 1H, NH).

**Example 33**

**N-[(3S)-1-Azabicyclo[2.2.2]oct-3-yl]-9H-xanthene-9-carboxamide**

The title compound was synthesised according to method a. The yield was 1.1 g, 60%.

20 m.p.: 244.2-244.9°C.

MS [M+1]<sup>+</sup>: 335.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 1.25-1.60 (m, 3H), 1.66 (m, 1H), 1.82 (m, 1H), 2.42-2.50 (m, 1H),  
2.55-2.75 (m, 3H), 2.83 (m, 1H), 3.05 (m, 1H), 3.60 (m, 1H), 5.0 (s, 1H), 7.05-7.15 (m,  
4H), 7.25-7.35 (m, 4H), 8.53 (d, 1H, NH).

25

**Example 34**

**(3S)-1-Allyl-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane trifluoroacetate**

The title compound was synthesised according to methods a and e. The yield of the final  
30 step was 0.0091 g, 22.1%.

MS [M-CF<sub>3</sub>COO]<sup>+</sup>: 375.

**Example 35**

**(3S)-1-Heptyl-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane trifluoroacetate**  
35



The title compound was synthesised according to methods a and e. The yield of the final step was 0.0097 g, 21.0 %.

MS [M-CF<sub>3</sub>COO]<sup>+</sup>: 433.

5 **Example 36**

**(3S)-1-Cyclohexylmethyl-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane trifluoroacetate**

The title compound was synthesised according to methods a and e. The yield of the final step was 0.0124 g, 27.0 %.

10 MS [M-CF<sub>3</sub>COO]<sup>+</sup>: 431.

**Example 37**

**(3S)-1-(3-Cyclohexylpropyl)-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane trifluoroacetate**

15 The title compound was synthesised according to methods a and e. The yield of the final step was 0.0045 g, 9.3%.

MS [M-CF<sub>3</sub>COO]<sup>+</sup>: 459.

**Example 38**

20 **(3S)-1-(3-Phenoxypropyl)-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane bromide**

The title compound was synthesised according to methods a and d. The yield of the final step was 0.28 g, 85%.

m.p.: 279.0-280.4°C.

25 MS [M-Br]<sup>+</sup>: 469.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 1.80-2.0 (m, 3H), 2.05-2.30 (m, 4H), 3.20-3.30 (m, 1H), 3.30-3.65 (m, 6H), 3.82 (m, 1H), 4.05 (t, 2H), 4.11 (m, 1H), 5.10 (s, 1H), 6.90-7.0 (m, 3H), 7.05-7.20 (m, 4H), 7.25-7.42 (m, 6H), 9.10 (d, 1H, NH).

30 **Example 39**

**(3S)-1-[3-(5,6,7,8-Tetrahydronaphthalen-2-yloxy)propyl]-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane trifluoroacetate**

The title compound was synthesised according to methods a and e. The yield of the final step was 0.0114 g, 21.2%.

35 MS [M-CF<sub>3</sub>COO]<sup>+</sup>: 523.

**Example 40**

**(3S)-1-[5-(2,6-Dimethylphenoxy)pentyl]-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane trifluoroacetate**

- 5 The title compound was synthesised according to methods a and e. The yield of the final step was 0.0101 g, 18.7%.

MS [M-CF<sub>3</sub>COO]<sup>+</sup>: 525.

**Example 41**

- 10 **N-(1-Azabicyclo[2.2.2]oct-3-yl)-9-hydroxy-9H-fluorene-9-carboxamide**

The title compound was synthesised according to method b. The yield was 1.06 g, 17.6%.

m.p.: 230°C.

MS [M+1]<sup>+</sup>: 335.

- 15 <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.0-1.30 (m, 2H), 1.52 (m, 2H), 1.79 (m, 1H), 2.04 (m, 1H), 2.36 (m, 1H), 2.57 (m, 3H), 3.05 (m, 1H), 3.77 (m, 1H), 5.45 (bs, 1H), 5.71 (d, 1H, NH), 7.20-7.50 (m, 6H), 7.55-7.70 (m, 2H).

**Example 42**

- 20 **3-[[[(9-Hydroxy-9H-fluoren-9-yl)carbonyl]amino]-1-methyl-1-azoniabicyclo[2.2.2]octane bromide**

The title compound was synthesised according to methods b and d. The yield of the final step was 0.26 g, 83.8%.

m.p.: 197.5-203.6°C.

- 25 MS [M-Br]<sup>+</sup>: 349.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 1.80-2.0 (m, 3H), 2.10-2.25 (m, 2H), 2.96 (s, 3H), 3.35-3.70 (m, 5H), 3.78 (m, 1H), 4.15 (m, 1H), 6.82 (s, 1H, OH), 7.30 (m, 2H), 7.43 (m, 4H), 7.79 (m, 2H), 8.81 (d, 1H, NH).

- 30 **Example 43**

**3-[[[(9-Hydroxy-9H-fluoren-9-yl)carbonyl]amino]-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane bromide**

The title compound was synthesised according to methods b and d. The yield of the final step was 0.35 g, 87.5%.

- 35 m.p.: 264°C.

MS [M-Br]<sup>+</sup>: 469.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 1.80-2.0 (m, 3H), 2.10-2.30 (m, 4H), 3.30-3.55 (m, 5H), 3.67 (m, 2H), 3.79 (m, 1H), 4.08 (t, 2H), 4.20 (m, 1H), 6.82 (s, 1H, OH), 6.97 (m, 3H), 7.32 (m, 4H), 7.43 (m, 4H), 7.79 (m, 2H), 8.88 (d, 1H, NH).

5

#### Example 44

##### N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-9-hydroxy-9H-fluorene-9-carboxamide

The title compound was synthesized according to method b. The yield was 0.304g, 4%.

MS [M+1]<sup>+</sup>: 335

10 <sup>1</sup>H-RMN (CDCl<sub>3</sub>): δ 7.67 - 7.62 (m, 2H), 7.42 - 7.36 (m, 4H), 7.32 - 7.25 (m, 2H), 6.08 - 5.93 (m, NH, 1H), 3.74 (m, 1H), 3.00 - 2.87 (m, 1H), 2.52 - 2.27 (m, 4H), 2.02 - 1.92 (m, 1H), 1.80 - 1.76 (m, 1H), 1.51 - 1.44 (m, 2H), 1.26 - 1.17 (m, 2H).

#### Example 45

##### 15 N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-fur-2-yl-2-hydroxypent-3-ynamide

The title compound was synthesized as a mixture of diastereomers according to method c. The yield was 1.38 g, 39%

MS [M+1]<sup>+</sup>: 289

20 <sup>1</sup>H-RMN (CDCl<sub>3</sub>): (mixture of diastereomers) δ 7.36 - 7.35 (m, 1H), 6.87 - 6.76 (m, NH, 1H), 6.54 - 6.52 (m, 1H), 6.34 - 6.32 (m, 1H), 3.92 (m, 1H), 3.29 - 3.17 (m, 1H), 2.78 - 2.64 (m, 4H), 2.47 - 2.35 (m, 1H), 2.00 - 1.90 (m, 1H), 1.92 (s, 3H), 1.70 - 1.39 (m, 4H).

#### Example 46

##### 25 N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-fur-2-yl-2-hydroxy-4-(4-methoxyphenyl)butanamide

The title compound was prepared as a mixture of diastereomers according to method c.

The yield was 2.4 g, 44%

MS [M+1]<sup>+</sup>: 385

30 <sup>1</sup>H-RMN (CDCl<sub>3</sub>): (mixture of diastereomers) δ 7.39 - 7.38 (m, 1H), 7.09 (d, 2H), 6.82 (d, 2H), 6.70 - 6.67 (m, NH, 1H), 6.41 - 6.35 (m, 2H), 3.92 (m, 1H), 3.78 (s, 3H), 3.33 - 3.24 (m, 1H), 2.82 - 2.19 (m, 9H), 1.96 - 1.87 (m, 1H), 1.67 - 1.40 (m, 4H)

#### Example 47

##### N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-N-methyl-9H-xanthene-9-carboxamide

35 The title product was synthesized according to method a. The yield was 0.9 g, 50%.

MS [M+1]<sup>+</sup>: 349

<sup>1</sup>H-RMN (400 MHz, 60°C, CDCl<sub>3</sub>): δ 7.26 - 7.19 (m, 4H), 7.10 - 7.01 (m, 4H), 5.51 (s, 1H), 4.18 (broad multiplet, 1H), 2.92 (s, 3H), 2.85-2.40 (m, 6H), 1.85-1.20 (m, 5H)

5 **Example 48**

**N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-(5-bromothiophen-2-yl)-2-(4-fluoro-3-methylphenyl)-2-hydroxyacetamide**

The title compound was synthesized as a mixture of diastereomers according to method c. The yield was 2.1 g, 42%.

10 MS [M+1]<sup>+</sup>: 453, 455

<sup>1</sup>H-RMN (CDCl<sub>3</sub>): (mixture of diastereomers) δ 7.35 - 7.20 (m, 2H), 7.08 - 6.89 (m, 3H), 6.80 (dd, 1H), 3.93 - 3.84 (m, 1H), 3.18 - 3.07 (m, 1H), 2.69 - 2.58 (m, 4H), 2.33 - 2.22 (m, 4H), 1.92 - 1.87 (m, 1H), 1.63 - 1.42 (m, 4H)

15 **Example 49**

**(2S)-N-[(3R)-1-azabicyclo[2.2.2]oct-3-yl]-2-cyclopentyl-2-hydroxy-2-thien-2-ylacetamide**

The title compound was synthesized according to method b. The yield was 0.650 g, 44%.

MS [M+1]<sup>+</sup>: 335

20 <sup>1</sup>H-RMN (CDCl<sub>3</sub>): δ 7.24 (m, 1H), 7.08 (m, 1H), 6.97 (m, 1H), 6.64 (d, NH, 1H), 3.91 - 3.81 (m, 1H), 3.33 - 3.23 (m, 1H), 2.88 - 2.69 (m, 5H), 2.42 - 2.33 (m, 1H), 1.87 (m, 1H), 1.69 - 1.36 (m, 12H)

**Example 50**

25 **(3R)-1-[3-(Benzothiazol-2-yloxy)propyl]-3-(2-fur-2-yl-2-hydroxypent-3-ynoylamino)-1-azonia-bicyclo[2.2.2]octane chloride**

The title compound was synthesized as a mixture of diastereomers according to methods c and d.

The reaction time for the final step (conditions: THF, reflux temperature) was 7 days. The obtained product was purified by several washings with ether. The yield was 0.337 g (75%).

30

MS [M-Cl]<sup>+</sup>: 480

HPLC: mixture of diastereomers 49:51

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): (mixture of diastereomers) δ 8.53 - 8.49 (m, NH, 1H), 7.69 (m, 1H), 7.58 (m, 1H), 7.53 - 7.39 (m, 2H), 7.26 - 7.19 (m, 2H), 6.43 - 6.38 (m, 2H), 4.20 - 4.14 (m, 1H), 4.05 - 3.99 (m, 2H), 3.76 - 3.65 (m, 1H), 3.60-3.20 (m, 7H), 2.20-1.65 (m, 7H), 1.87 (s, 3H)

5

**Example 51**

**(3R)-3-(2-Fur-2-yl-2-hydroxypent-3-ynoylamino)-1-[3-(naphthalen-1-yloxy)propyl]-1-azoniabicyclo[2.2.2]octane chloride**

The title compound was synthesized as a mixture of diastereomers according to methods c and d.

The reaction time for the final step (conditions, THF, reflux temperature) was 7 days. The obtained product was purified by several washings with ether. The yield was 0.252 g (57%).

MS [M-Cl]<sup>+</sup>: 473

15 HPLC: mixture of diastereomers 48:52

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): (mixture of diastereomers) δ 8.62 - 8.58 (m, NH, 1H), 8.23 (m, 1H), 7.90 - 7.86 (m, 1H), 7.61 - 7.40 (m, 5H), 7.25 (m, 1H), 6.99 (m, 1H), 6.46 - 6.39 (m, 2H), 4.30 - 4.20 (m, 3H), 3.95 - 3.30 (m, 8H), 2.42 - 2.25 (m, 2H), 2.20 - 1.70 (m, 5H), 1.89 (s, 3H).

20

**Example 52**

**(3R)-1-[3-(Benzo[1,3]dioxol-5-yloxy)propyl]-3-(2-fur-2-yl-2-hydroxypent-3-ynoylamino)-1-azonia-bicyclo[2.2.2]octane bromide**

The title compound was synthesised as a mixture of diastereomers according to methods c and d.

The reaction time for the final step (conditions, THF, reflux temperature) was 2 days. The obtained product was purified by several washings with ether. The yield was 0.320 g (78%).

MS [M-Br]<sup>+</sup>: 467

30 HPLC: mixture of diastereomers 52:48

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): (mixture of diastereomers) δ 8.54 - 8.49 (m, NH, 1H), 7.60 (s, 1H), 7.21 (d, 1H), 6.82 (d, 1H), 6.65 (m, 1H), 6.45 - 6.36 (m, 3H), 5.96 (s, 2H), 4.26 - 4.15 (m, 1H), 4.00 - 3.95 (m, 2H), 3.82 - 3.70 (m, 1H), 3.61 - 3.28 (m, 7H), 2.15 - 1.74 (m, 7H), 1.89 (s, 3H).

35

**Example 53**

**(3R)-3-[[[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetyl]amino]-1-(4,4,4-trifluorobutyl)-1-azoniabicyclo[2.2.2]octane bromide**

The title compound was synthesised according to method d from Example 49.

- 5 The reaction time (conditions, THF, reflux temperature) was 3 days. The obtained product was purified by several washings with ether. The yield was 0.183 g (77%).

MS [M-Br]<sup>+</sup>: 445

- <sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 8.28 (d, NH, 1H), 7.38 (dd, 1H), 7.09 (dd, 1H), 6.94 (dd, 1H), 6.11 (s, OH, 1H), 4.21 - 4.12 (m, 1H), 3.72 (m, 1H), 3.59 - 3.30 (m, 5H), 3.25 - 3.17 (m, 2H), 2.80 (m, 1H), 2.41 - 2.25 (m, 2H), 2.05 - 1.80 (m, 6H), 1.80-1.62 (m, 1H), 1.62-1.30 (m, 8H)

**Example 54**

- 15 **(3R)-3-[[[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetyl]amino]-1-(2-hydroxyethyl)-1-azoniabicyclo[2.2.2]octane bromide**

The title compound was synthesised according to method d from Example 49.

The reaction time (conditions: THF, reflux temperature) was 5 days. The obtained product was purified by several washings with ether. The yield was 0.174 g (84%).

MS [M-Br]<sup>+</sup>: 379

- 20 <sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 8.24 (d, NH, 1H), 7.37 (dd, 1H), 7.08 (dd, 1H), 6.94 (dd, 1H), 6.10 (s, OH, 1H), 5.27 (t, OH, 1H), 5.20-5.10 (m, 1H), 4.15 (m, 1H), 3.85 - 3.75 (m, 2H), 3.75-3.20 (m, 5H), 2.83 - 2.77 (m, 1H), 2.00 - 1.85 (m, 4H), 1.77 - 1.66 (m, 1H), 1.53 - 1.41 (m, 8H)

25 **Example 55**

**(3R)-1-(4-Acetoxybutyl)-3-[2-(5-bromothien-2-yl)-2-(4-fluoro-3-methylphenyl)-2-hydroxyacetyl]amino]-1-azoniabicyclo[2.2.2]octane bromide**

- 30 The title compound was synthesised as a mixture of diastereomers according to methods c and d. The reaction time for the final step (conditions: THF, reflux temperature) was 41 hours. The obtained product was purified by several washings with ether. The yield was 0.190 g (88%).

MS [M-Br]<sup>+</sup>: 569, 567

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): (mixture of diastereomers) δ 8.72 (d, NH, 1H), 7.45 (d, 1H), 7.36 - 7.18 (m, 2H), 7.13 - 7.06 (m, 2H), 6.91 (t, 1H), 4.25 - 4.18 (m, 1H), 4.06 - 4.00 (m, 2H),

3.72 - 3.26 (m, 6H), 3.19 - 3.11 (m, 2H), 2.20 (s, 3H), 2.21 - 1.84 (m, 1H), 2.02 (s, 3H), 1.95 - 1.54 (m, 8H)

#### Example 56

5 **(3R)-3-[2-(5-Bromothiophen-2-yl)-2-(4-fluoro-3-methylphenyl)-2-hydroxyacetyl-amino]-1-(4-ethoxycarbonylbutyl)-1-azoniabicyclo[2.2.2]octane bromide**

The title compound was synthesised as a mixture of diastereomers according to methods c and d. The reaction time for the final step (conditions: THF, reflux temperature) was 46 hours. The obtained product was purified by several washings with ether. The yield was

10 0.201 g (92%).

MS [M-Br]<sup>+</sup>: 583, 581

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): (mixture of diastereomers) δ 8.72 (d, NH, 1H), 7.45 (d, 1H), 7.36 - 7.18 (m, 2H), 7.13 - 7.06 (m, 2H), 6.91 (t, 1H), 4.26 - 4.16 (m, 1H), 4.06 (q, 2H), 3.65 - 3.30 (m, 6H), 3.18 - 3.09 (m, 2H), 2.40 - 2.32 (m, 2H), 2.20 (s, 3H), 2.08 - 1.48 (m, 9H),

15 1.18 (t, 3H)

#### Example 57

**(3R)-1-(3-Cyanopropyl)-3-[2-fur-2-yl-2-hydroxy-4-(4-methoxyphenyl)butyrylamino]-1-azoniabicyclo[2.2.2]octane bromide**

20 The title compound was synthesised as a mixture of diastereomers according to methods c and d. The reaction time for the final step (conditions: THF, reflux temperature) was 3 days. The obtained product was purified by several washings with ether. The yield was 0.113 g (65%).

MS [M-Br]<sup>+</sup>: 452

25 <sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): (mixture of diastereomers) δ 8.49 - 8.40 (m, NH, 1H), 7.58 (s, 1H), 7.09 (d, 2H), 6.84 (d, 2H), 6.41 - 6.35 (m, 2H), 6.33 (d, 1H), 4.27 - 4.19 (m, 1H), 3.80-3.65 (m, 1H), 3.71 (s, 3H), 3.63 - 3.36 (m, 5H), 3.23 - 3.17 (m, 2H), 2.68 - 2.60 (m, 3H), 2.40 - 1.73 (m, 10H)

#### 30 Example 58

**(3R)-1-(2-[1,3]Dioxolan-2-yl-ethyl)-3-[2-fur-2-yl-2-hydroxy-4-(4-methoxyphenyl)butyrylamino]-1-azoniabicyclo[2.2.2]octane bromide**

The title compound was synthesised as a mixture of diastereomers according to methods c and d. The reaction time for the final step (conditions: THF, reflux temperature) was 3

days. The obtained product was purified by several washings with ether. The yield was 0.128 g (70%).

MS [M-Br]<sup>+</sup>: 485

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): (mixture of diastereomers) δ 8.47 - 8.37 (m, NH, 1H), 7.58 (s, 1H),  
5 7.09 (d, 2H), 6.84 (d, 2H), 6.40 - 6.35 (m, 2H), 6.32 (d, 1H), 4.94 - 4.90 (m, 1H), 4.21 (m,  
1H), 3.96 - 3.77 (m, 4H), 3.71 (s, 3H), 3.59 - 3.21 (m, 7H), 2.72 - 2.61 (m, 1H), 2.41 - 1.74  
(m, 10H).

#### Example 59

10 **(3R)-3-[(9-Hydroxy-9H-fluoren-9-yl)carbonyl]amino]-1-[3-(methylphenylamino)propyl]-1-azoniabicyclo[2.2.2]octane chloride**

The title compound was synthesised according to methods b and d. The reaction time for the final step (conditions: THF:CHCl<sub>3</sub> 1:1, reflux temperature) was 20 days. The obtained product was purified by washing with hot CHCl<sub>3</sub> and ether. The yield was 0.052 g (33%).

15 MS [M-Cl]<sup>+</sup>: 482

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 8.86 (d, NH, 1H), 7.78 (d, 2H), 7.45 - 7.16 (m, 8H), 6.81 - 6.62 (m,  
, 4H), 4.23 - 4.14 (m, 1H), 3.76 - 3.53 (m, 2H), 3.43 - 3.17 (m, 8H), 2.91 (s, 3H), 2.23 -  
1.77 (m, 7H).

#### 20 Example 60

**(3R)-3-[Methyl-(9H-xanthen-9-yl)carbonyl]amino]-1-(3-pyrrol-1-ylpropyl)-1-azoniabicyclo[2.2.2]octane bromide**

The title compound was synthesised according to methods a and d. The reaction time for the final step (conditions: THF, reflux temperature) was 2 days. The obtained product was  
25 purified by several washings with THF and ether. The yield was 0.257 g (83%).

MS [M-Br]<sup>+</sup>: 456

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 7.34 - 7.06 (m, 8H), 6.78 (m, 2H), 6.01 (m, 2H), 5.75 (s, 1H),  
4.47 (m, 1H), 3.94 - 3.88 (m, 2H), 3.72 - 3.25 (m, 6H), 3.39 (s, 3H), 3.20-3.05 (m, 2H),  
2.22 - 1.88 (m, 7H).

30

#### Example 61

**(3R)-1-[3-(Biphenyl-4-yloxy)propyl]-3-[methyl-(9H-xanthene-9-carbonyl)amino]-1-azoniabicyclo[2.2.2]octane chloride**



The title compound was synthesised according to methods a and d. The reaction time for the final step (conditions: THF, reflux temperature) was 11 days. The obtained product was purified by several washings with THF and ether. The yield was 0.195 g (57%).

MS [M-Cl]<sup>+</sup>: 559

- 5 <sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 7.62 (d, 4H), 7.47 - 7.02 (m, 13H), 5.77 (s, 1H), 4.50 (m, 1H), 4.13 - 4.06 (m, 2H), 3.82 - 3.31 (m, 8H), 3.42 (s, 3H), 2.25 - 1.91 (m, 7H).

The following examples illustrate pharmaceutical compositions according to the present invention and procedures for their preparation.

### Example 62

#### Preparation of a pharmaceutical composition: tablets

##### Formulation:

15	Compound of the present invention	5.0 mg
	Lactose	113.6 mg
	Microcrystalline cellulose	28.4 mg
	Light silicic anhydride	1.5 mg
	Magnesium stearate	1.5 mg

20

Using a mixer machine, 15 g of the compound of the present invention was mixed with 340.8 g of lactose and 85.2 g of microcrystalline cellulose. The mixture was subjected to compression moulding using a roller compactor to give a flake-like compressed material. The flake-like compressed material was pulverized using a hammer mill, and the

25 pulverized material was screened through a 20 mesh screen. A 4.5 g portion of light silicic anhydride and 4.5 g of magnesium stearate were added to the screened material and mixed. The mixer product was subjected to a tablets making machine equipped with a die/punch system of 7.5 mm in diameter, thereby obtaining 3,000 tablets each having 150 mg in weight.

30

### Example 63

#### Preparation of a pharmaceutical composition: coated tablets

##### Formulation:

	Compound of the present invention	5.0 mg
35	Lactose	95.2 mg

	Corn starch	40.8 mg
	Polyvinylpyrrolidone K25	7.5 mg
	Magnesium stearate	1.5 mg
	Hydroxypropylcellulose	2.3 mg
5	Polyethylene glycol 6000	0.4 mg
	Titanium dioxide	1.1 mg
	Purified talc	0.7 mg

Using a fluidized bed granulating machine, 15 g of the compound of the present invention  
10 was mixed with 285.6 g of lactose and 122.4 g of corn starch. Separately, 22.5 g of  
polyvinylpyrrolidone was dissolved in 127.5 g of water to prepare a binding solution. Using  
a fluidized bed granulating machine, the binding solution was sprayed on the above  
mixture to give granulates. A 4.5 g portion of magnesium stearate was added to the  
obtained granulates and mixed. The obtained mixture was subjected to a tablet making  
15 machine equipped with a die/punch biconcave system of 6.5 mm in diameter, thereby  
obtaining 3,000 tablets, each having 150 mg in weight.

Separately, a coating solution was prepared by suspending 6.9 g of  
hydroxypropylmethylcellulose 2910, 1.2 g of polyethylene glycol 6000, 3.3 g of titanium  
dioxide and 2.1 g of purified talc in 72.6 g of water. Using a High Coated, the 3,000 tablets  
20 prepared above were coated with the coating solution to give film-coated tablets, each  
having 154.5 mg in weight.

#### Example 64

##### Preparation of a pharmaceutical composition: liquid inhalant

25 Formulation:

Compound of the present invention	400 $\mu$ g	
Physiological saline		1 ml

A 40 mg portion of the compound of the present invention was dissolved in 90 ml of  
30 physiological saline, and the solution was adjusted to a total volume of 100 ml with the  
same saline solution, dispensed in 1 ml portions into 1 ml capacity ampoule and then  
sterilized at 115 °C for 30 minutes to give a liquid inhalant.

#### Example 65

35 Preparation of a pharmaceutical composition: powder inhalant

**Formulation:**

Compound of the present invention	200 µg
Lactose	4,000 µg

- 5 A 20 g portion of the compound of the present invention was uniformly mixed with 400 g of lactose, and a 200 mg portion of the mixture was packed in a powder inhaler for exclusive use to produce a powder inhalant.

**Example 66**

10 **Preparation of a pharmaceutical composition: inhalation aerosol.**

**Formulation:**


Compound of the present invention	200 µg
Dehydrated (Absolute) ethyl alcohol USP	8,400 µg
1,1,1,2-Tetrafluoroethane (HFC-134A)	46,810 µg

15

- The active ingredient concentrate is prepared by dissolving 0.0480 g of the compound of the present invention in 2.0160 g of ethyl alcohol. The concentrate is added to an appropriate filling apparatus. The active ingredient concentrate is dispensed into aerosol container, the headspace of the container is purged with Nitrogen or HFC-134A vapour
- 20 (purging ingredients should not contain more than 1 ppm oxygen) and is sealed with valve. 11.2344 g of HFC-134A propellant is then pressure filled into the sealed container.

CLAIMS

1. A compound which is a quinuclidine amide derivative of formula (I):




(I)

wherein


$R_1$  represents a hydrogen atom or an straight or branched, optionally substituted lower alkyl group;

$R_2$  represents a group of formula i) or ii)

i)



ii)



wherein

$R_3$  represents a group selected from phenyl, 2-furyl, 3-furyl, 2-thienyl or 3-thienyl;

$R_4$  represents a group selected from optionally substituted lower alkyl, optionally substituted lower alkenyl, optionally substituted lower alkynyl, cycloalkyl, cycloalkylmethyl, phenyl, benzyl, phenethyl, 2-furyl, 3-furyl, 2-thienyl or 3-thienyl;

and  $R_5$  represents a hydrogen atom or a hydroxy, methyl, or  $-CH_2OH$  group;

the benzene rings in formula ii) and the cyclic groups represented by  $R_3$  and  $R_4$  being each independently optionally substituted by one, two or three substituents selected from

halogen, straight or branched, optionally substituted lower alkyl, hydroxy, straight or branched, optionally substituted lower alkoxy, nitro, cyano,  $-\text{CO}_2\text{R}'$  or  $-\text{NR}'\text{R}''$ , wherein  $\text{R}'$  and  $\text{R}''$  each independently represents a hydrogen atom or a straight or branched, optionally substituted lower alkyl group or  $\text{R}'$  and  $\text{R}''$  together with the atom to which they are attached form a cyclic group;

$\text{Q}$  represents a single bond or a  $-\text{CH}_2-$ ,  $-\text{CH}_2\text{-CH}_2-$ ,  $-\text{O}-$ ,  $-\text{O-CH}_2-$ ,  $-\text{S}-$ ,  $-\text{S-CH}_2-$  or  $-\text{CH=CH}-$  group;

$p$  is 1 or 2 and the amide group is at positions 2, 3 or 4 of the azabicyclic ring;

or pharmaceutically acceptable salts thereof, including quaternary ammonium salts;

and all individual stereoisomers and mixtures thereof;

15


with the proviso that when  $p$  is 2, the amide moiety is in position 3 of the quinuclidine ring,  $\text{R}_1$  is hydrogen and  $\text{R}_3$  and  $\text{R}_4$  are both unsubstituted phenyl, then

-when said compound is not a pharmaceutically acceptable salt or is a HCl salt, then  $\text{R}_5$

20 cannot be one of hydrogen or hydroxy; and

- when said compound is a quaternary ammonium salt having a methyl group attached to the nitrogen atom of the quinuclidine ring, then  $\text{R}_5$  cannot be hydroxy.

2. A compound according to claim 1 which is a quaternary ammonium salt of formula (II)



(II)

25

wherein  $\text{R}_1$ ,  $\text{R}_2$  and  $p$  are as defined in claim 1;

$m$  is an integer from 0 to 8;

30

$n$  is an integer from 0 to 4;

A represents a group selected from  $-\text{CH}_2-$ ,  $-\text{CH}=\text{CR}'-$ ,  $-\text{CR}'=\text{CH}-$ ,  $-\text{CR}'\text{R}''-$ ,  $-\text{C}(\text{O})-$ ,  $-\text{O}-$ ,  $-\text{S}-$ ,  $-\text{S}(\text{O})-$ ,  $-\text{S}(\text{O})_2-$  and  $-\text{NR}'-$ , wherein  $\text{R}'$  and  $\text{R}''$  are as defined in claim 1;

- 5  $\text{R}_6$  represents a hydrogen atom, or a group selected from straight or branched, optionally substituted lower alkyl, hydroxy, straight or branched, optionally substituted lower alkoxy, cyano, nitro,  $-\text{CH}=\text{CR}'\text{R}''$ ,  $-\text{C}(\text{O})\text{OR}'$ ,  $-\text{OC}(\text{O})\text{R}'$ ,  $-\text{SC}(\text{O})\text{R}'$ ,  $-\text{C}(\text{O})\text{NR}'\text{R}''$ ,  $-\text{NR}'\text{C}(\text{O})\text{OR}''$ ,  $-\text{NR}'\text{C}(\text{O})\text{NR}''$ , cycloalkyl, phenyl, naphthanelyl, 5,6,7,8-tetrahydronaphthanelyl, benzo[1,3]dioxolyl, heteroaryl or heterocyclyl;  $\text{R}'$  and  $\text{R}''$  being as defined in claim 1; and
- 10 wherein the cyclic groups represented by  $\text{R}_6$  are optionally substituted by one, two or three substituents selected from halogen, hydroxy, straight or branched, optionally substituted lower alkyl, phenyl,  $-\text{OR}'$ ,  $-\text{SR}'$ ,  $-\text{NR}'\text{R}''$ ,  $-\text{NHCOR}'$ ,  $-\text{CONR}'\text{R}''$ ,  $-\text{CN}$ ,  $-\text{NO}_2$  and  $-\text{COOR}'$ ;  $\text{R}'$  and  $\text{R}''$  being as defined in claim 1; and
- 15  $\text{X}^-$  represents a pharmaceutically acceptable anion of a mono or polyvalent acid.

and all individual stereoisomers and mixtures thereof;

with the proviso that when  $p$  is 2, the amide moiety is in position 3 of the quinuclidine ring,

- 20  $\text{R}_1$  is hydrogen,  $\text{R}_3$  and  $\text{R}_4$  are both unsubstituted phenyl and  $\text{R}_5$  is hydroxy, then in the compounds of formula (II) the sequence  $\text{R}_6 - (\text{CH}_2)_n - \text{A} - (\text{CH}_2)_m -$  cannot be a methyl group.

3. A compound according to claim 1 or claim 2 wherein  $\text{R}_1$  is hydrogen, methyl or ethyl.

25

4. A compound according to claim 3, wherein  $\text{R}_1$  is hydrogen.

5. A compound according to any one of the preceding claims, wherein  $\text{R}_2$  is a group of formula i), wherein  $\text{R}_3$  is a group, which is optionally substituted with one or more halogen
- 30 atom(s), selected from phenyl, 2-thienyl, 3-thienyl or 2-furyl

6. A compound according to claim 5, wherein  $\text{R}_2$  is a group of formula i), wherein  $\text{R}_3$  represents a group phenyl, 2-thienyl or 2-furyl which are optionally substituted with one or more halogen atom(s).

35

7. A compound according to any one of claims 5 and 6, wherein  $R_2$  is a group of formula i), wherein  $R_3$  represents phenyl or 2-thienyl.

5 8. A compound according to any one of the preceding claims, wherein  $R_4$  represents a linear group selected from ethyl, n-butyl, vinyl, allyl, 1-propenyl and 1-propynyl, or a group, which is optionally substituted with one or more halogen atom(s), methyl or methoxy group(s), selected from cyclopentyl, cyclohexyl, phenyl, benzyl, phenethyl, 2-thienyl and 3-furyl.

10

9. A compound according to claim 8, wherein  $R_4$  represents a linear group selected from ethyl, n-butyl, vinyl, allyl and 1-propynyl, or a group, which is optionally substituted with one or more halogen atom(s), methyl or methoxy group(s), selected from cyclopentyl, phenyl, benzyl, phenethyl and 2-thienyl.

15

10. A compound according to any one of claims 8 and 9, wherein  $R_4$  represents a group selected from ethyl, n-butyl, vinyl, allyl, cyclopentyl, phenyl, benzyl or 2-thienyl

20

11. A compound according to any one of claims 1 to 4, wherein  $R_2$  is a group of formula ii), wherein Q represents a single bond or an oxygen atom.

12. A compound according to any one of the preceding claims wherein  $R_5$  is hydrogen or hydroxy.

25

13. A compound according to any one of the preceding claims wherein p is 2 and the amide group is at positions 3 or 4 of the azabicyclic ring.

14. A compound according to claim 13 wherein the amide group is at position 3 of the azabicyclic ring.

30

15. A compound of formula (II) according to any one of claims 2 to 14, wherein m is an integer from 0 to 6 and n is an integer from 0 to 4; A represents a group selected from  $-CH_2-$ ,  $-CH=CH-$ ,  $-O-$ ,  $-C(O)-$ ,  $-NR'-$ , and  $-S-$ ; and  $R_6$  is a hydrogen atom, a cyano group, a nitro group, a  $-C(O)OR'$  group, a  $-OC(O)R'$  group, a  $-SC(O)R'$  group, a  $-CH=CH_2$  group, 35 a  $-CH=CR'R''$  group, a  $C(O)NR'R''$  group, a straight or branched  $C_1-C_4$  alkyl group, which

is optionally substituted with one or more halogen atom(s), a straight C<sub>1</sub>-C<sub>4</sub> alkoxy group, which is optionally substituted with one or more halogen atom(s) or hydroxy group(s), or a cyclic group, which is optionally substituted with one or more substituents selected from halogen atoms, groups of formula -C(O)NR'R'' and methyl, hydroxy, nitro and phenyl groups, the cyclic group being selected from cyclohexyl, phenyl, 5,6,7,8-tetrahydronaphthalenyl, 2-thienyl, 1-pyrrolidinyl, 1-pyrrolyl, benzo[1,3]dioxolyl, 2-benzothiazolyl, naphthalenyl and dioxolyl.

16. A compound according to claim 15, wherein m is an integer from 0 to 5 and n is an integer from 0 to 2; A represents a group selected from -CH<sub>2</sub>-, -CH=CH-, -O-, -C(O)-, -NR'-, and -S-; and R<sub>6</sub> is a hydrogen atom, a cyano group, a -C(O)OR' group, a -OC(O)R' group, a -SC(O)R' group, a -CH=CH<sub>2</sub> group, a -C(O)NR'R'' group, a straight or branched C<sub>1</sub>-C<sub>4</sub> alkyl group, a trifluoromethyl, or a cyclic group selected from cyclohexyl, 5,6,7,8-tetrahydronaphthalenyl, 2-thienyl, 1-pyrrolyl, benzo[1,3]dioxolyl, 2-benzothiazolyl, naphthalenyl, dioxolyl and phenyl, which is optionally substituted with one or more substituents selected from halogen atoms, groups of formula -C(O)NR'R'', methyl, hydroxy and phenyl groups.

17. A compound according to any one of claims 15 and 16, wherein m is an integer from 0 to 5 and n is an integer from 0 to 2; A represents a group selected from -CH<sub>2</sub>-, -CH=CH-, -O-; and R<sub>6</sub> is selected from hydrogen, straight C<sub>1</sub>-C<sub>4</sub> alkyl, -CH=CH<sub>2</sub>, cyclohexyl, phenyl which is unsubstituted or substituted with one or two substituents selected from methyl groups and hydroxy groups, 5,6,7,8-tetrahydronaphthalenyl and 2-thienyl.

18. A compound according to claim 17, wherein the sequence R<sub>6</sub> - (CH<sub>2</sub>)<sub>n</sub> - A - (CH<sub>2</sub>)<sub>m</sub> - is one of methyl, 3-phenoxypropyl, 3-(3-hydroxyphenoxy)propyl, allyl, heptyl, 3-phenylpropyl, 3-phenylallyl, 2-phenoxyethyl, 2-benzyloxyethyl, cyclohexylmethyl, 3-(5,6,7,8-tetrahydronaphthalen-2-yloxy)propyl, 5-(2,6-dimethylphenoxy)pentyl, 3-thien-2-ylpropyl or 3-cyclohexylpropyl and X<sup>-</sup> is bromide or trifluoroacetate.

19. A compound according to any one of the preceding claims, which is a single isomer.

20. A compound according to claim 1 or claim 2 which is one of:

N-(1-Azabicyclo[2.2.2]oct-3-yl)-2-hydroxy-2,2-dithien-2-ylacetamide

N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2,2-dithien-2-ylacetamide



- N-[(3S)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2,2-dithien-2-ylacetamide  
N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2,2-dithien-2-ylacetamide  
N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-cyclopentyl-2-hydroxy-2-thien-2-ylacetamide  
N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2-thien-2-ylpent-4-enamide
- 5 (2\*)-N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2-thien-2-ylbutanamide  
(diastereomer 1)  
(2\*)-N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2-thien-2-ylbutanamide  
(diastereomer 2)  
(2\*)-N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2-thien-2-ylbut-3-enamide
- 10 (diastereomer 1)  
(2\*)-N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2-thien-2-ylbut-3-enamide  
(diastereomer 2)  
N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2,3-diphenylpropanamide  
N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-cyclopentyl-2-hydroxy-2-phenylacetamide
- 15 N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-phenylhexanamide  
N-(1-Azabicyclo[2.2.2]oct-3-yl)-9H-xanthene-9-carboxamide  
N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-9H-xanthene-9-carboxamide  
N-[(3S)-1-Azabicyclo[2.2.2]oct-3-yl]-9H-xanthene-9-carboxamide  
N-(1-Azabicyclo[2.2.2]oct-3-yl)-9-hydroxy-9H-fluorene-9-carboxamide
- 20 N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-N-methyl-9H-xanthene-9-carboxamide  
(2S)-N-[(3R)-1-azabicyclo[2.2.2]oct-3-yl]-2-cyclopentyl-2-hydroxy-2-thien-2-ylacetamide  
3-(2-Hydroxy-2,2-dithien-2-ylacetyl-amino)-1-methyl-1-azoniabicyclo[2.2.2]octane bromide  
3-(2-Hydroxy-2,2-dithien-2-ylacetyl-amino)-1-(3-phenoxypropyl)-1-  
azoniabicyclo[2.2.2]octane bromide
- 25 (3R)-3-(2-Hydroxy-2,2-dithien-2-ylacetyl-amino)-1-methyl-1-azoniabicyclo[2.2.2]octane  
trifluoroacetate  
(3R)-1-Allyl-3-(2-hydroxy-2,2-dithien-2-ylacetyl-amino)-1-azoniabicyclo[2.2.2]octane  
trifluoroacetate  
(3R)-1-Heptyl-3-(2-hydroxy-2,2-dithien-2-ylacetyl-amino)-1-azoniabicyclo[2.2.2]octane
- 30 trifluoroacetate  
(3R)-3-(2-Hydroxy-2,2-dithien-2-ylacetyl-amino)-1-(3-phenylpropyl)-1-  
azoniabicyclo[2.2.2]octane bromide  
(3R)-3-(2-Hydroxy-2,2-dithien-2-ylacetyl-amino)-1-((E)-3-phenylallyl)-1-  
azoniabicyclo[2.2.2]octane trifluoroacetate

- (3R)-3-(2-Hydroxy-2,2-dithien-2-ylacetyl amino)-1-(2-phenoxyethyl)-1-azoniabicyclo[2.2.2]octane bromide
- (3R)-3-(2-Hydroxy-2,2-dithien-2-ylacetyl amino)-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane bromide
- 5 (3R)-3-(2-Hydroxy-2,2-dithien-2-ylacetyl amino)-1-[3-(3-hydroxyphenoxy)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-1-(2-Benzoyloxyethyl)-3-(2-hydroxy-2,2-dithien-2-ylacetyl amino)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-(2-Hydroxy-2,2-dithien-2-ylacetyl amino)-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane bromide
- 10 (3S)-3-(2-Hydroxy-2,2-dithien-2-ylacetyl amino)-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane bromide
- (3R)-3-(2,2-Dithien-2-ylacetyl amino)-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane bromide
- 15 1-Methyl-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane bromide
- 1-(3-Phenoxypropyl)-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane bromide
- (3R)-1-(3-Phenoxypropyl)-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane bromide
- 20 (3S)-1-Allyl-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3S)-1-Heptyl-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3S)-1-Cyclohexylmethyl-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 25 (3S)-1-(3-Cyclohexylpropyl)-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3S)-1-(3-Phenoxypropyl)-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane bromide
- 30 (3S)-1-[3-(5,6,7,8-Tetrahydronaphthalen-2-yloxy)propyl]-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3S)-1-[5-(2,6-Dimethylphenoxy)pentyl]-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 35 3-[[[9-Hydroxy-9H-fluoren-9-yl)carbonyl]amino]-1-methyl-1-azoniabicyclo[2.2.2]octane bromide

- 3-[[[(9-Hydroxy-9H-fluoren-9-yl)carbonyl]amino]-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane bromide  
(3R)-1-[3-(2-Carbamoylphenoxy)propyl]-3-[[[(9-Hydroxy-9H-fluoren-9-yl)carbonyl]amino]-1-azoniabicyclo[2.2.2]octane formate
- 5 (3R)-1-[4-(4-Fluorophenyl)-4-oxobutyl]-3-[[[(9-Hydroxy-9H-fluoren-9-yl)carbonyl]amino]-1-azoniabicyclo[2.2.2]octane formate  
(3R)-3-[[[(9-Hydroxy-9H-fluoren-9-yl)carbonyl]amino]-1-[3-(methylphenylamino)propyl]-1-azoniabicyclo[2.2.2]octane chloride  
(3R)-3-[[[(9-Hydroxy-9H-fluoren-9-yl)carbonyl]amino]-1-(3-phenylsulfanylpropyl)-1-azoniabicyclo[2.2.2]octane formate
- 10 (3R)-3-[Methyl-(9H-xanthen-9-ylcarbonyl)amino]-1-(3-pyrrol-1-ylpropyl)-1-azoniabicyclo[2.2.2]octane bromide  
(3R)-1-[3-(Biphenyl-4-yloxy)propyl]-3-[methyl-(9H-xanthene-9-carbonyl)amino]-1-azoniabicyclo[2.2.2]octane chloride
- 15 (3R)-3-(2-Fur-2-yl-2-hydroxypent-3-ynoylamino)-1-[3-(naphthalen-1-yloxy)propyl]-1-azoniabicyclo[2.2.2]octane chloride  
(3R)-1-[3-(Benzo[1,3]dioxol-5-yloxy)propyl]-3-(2-fur-2-yl-2-hydroxypent-3-ynoylamino)-1-azonia-bicyclo[2.2.2]octane bromide  
(3R)-1-[3-(Benzothiazol-2-yloxy)propyl]-3-(2-fur-2-yl-2-hydroxypent-3-ynoylamino)-1-azonia-bicyclo[2.2.2]octane chloride
- 20 (3R)-3-[[[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetyl]amino]-1-(2-hydroxyethyl)-1-azoniabicyclo[2.2.2]octane bromide  
(3R)-3-[[[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetyl]amino]-1-(2-ethoxyethyl)-1-azoniabicyclo[2.2.2]octane formate
- 25 (3R)-3-[[[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetyl]amino]-1-(4,4,4-trifluorobutyl)-1-azoniabicyclo[2.2.2]octane bromide  
(3R)-1-(4-Acetoxybutyl)-3-[2-(5-bromothien-2-yl)-2-(4-fluoro-3-methylphenyl)-2-hydroxyacetyl-amino]-1-azoniabicyclo[2.2.2]octane bromide  
(3R)-3-[2-(5-Bromothien-2-yl)-2-(4-fluoro-3-methylphenyl)-2-hydroxyacetyl-amino]-1-(4-ethoxycarbonylbutyl)-1-azoniabicyclo[2.2.2]octane bromide
- 30 (3R)-1-(3-Acetylsulfanylpropyl)-3-[2-(5-bromothien-2-yl)-2-(4-fluoro-3-methylphenyl)-2-hydroxyacetyl-amino]-1-azoniabicyclo[2.2.2]octane formate  
(3R)-1-(3-Cyanopropyl)-3-[2-fur-2-yl-2-hydroxy-4-(4-methoxyphenyl)butyrylamino]-1-azoniabicyclo[2.2.2]octane bromide


(3R)-1-(2-Carbamoylethyl)-3-[2-fur-2-yl-2-hydroxy-4-(4-methoxyphenyl)butyrylamino]-1-azoniabicyclo[2.2.2]octane formate

(3R)-1-(2-[1,3]Dioxolan-2-yl-ethyl)-3-[2-fur-2-yl-2-hydroxy-4-(4-methoxyphenyl)butyrylamino]-1-azoniabicyclo[2.2.2]octane bromide

5

21. A process for producing a compound of formula (I), as defined in claim 1 and wherein  $R_2$  is a group of formula i) and  $R_5$  is an hydroxy group, which process comprises reacting a compound of formula (V).

10



(V)

wherein  $R_1$  and  $R_3$  are as defined in claim 1 with the corresponding organometallic derivative  $R_4$ -[Mg,Li], wherein  $R_4$  is as defined in claim 1.

22. A compound of formula (V), which is one of

15

N-1-azabicyclo[2.2.2]oct-3-yl-2-oxo-2-thien-2-ylacetamide

N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-oxo-2-thien-2-ylacetamide

N-[(3S)-1-Azabicyclo[2.2.2]oct-3-yl]-2-oxo-2-thien-2-ylacetamide

20 23. A process for producing a compound of formula (II), as defined in claim 2, which process comprises reacting a compound of formula (I), as defined in claim 1 with an alkylating agent of formula  $R_6$ -( $CH_2$ ) $_n$ -A-( $CH_2$ ) $_m$ -W, wherein  $R_6$ ,  $n$ , A and  $m$  are as defined in claim 2 and W represents any suitable leaving group.

25 24. A pharmaceutical composition comprising a compound according to any one of claims 1 to 20 in admixture with a pharmaceutically acceptable carrier or diluent.

25. A compound according to any one of claims 1 to 20 for the treatment of a pathological condition or disease susceptible to amelioration by antagonism of M3 muscarinic receptors.
- 5 26. Use of a compound according to any one of claims 1 to 20 in the manufacture of a medicament for the treatment of a pathological condition or disease susceptible to amelioration by antagonism of M3 muscarinic receptors.
27. Use according to claim 26 wherein the pathological condition is a respiratory,  
10 urological or gastrointestinal disease or disorder.
28. A method for treating a subject afflicted with a pathological condition or disease susceptible to amelioration by antagonism of M3 muscarinic receptors, which comprises administering to said subject an effective amount of a compound as defined in any one of  
15 claims 1 to 20.
29. A method according to claim 28 wherein the pathological condition is a respiratory, urological or gastrointestinal disease or disorder.
- 20 30. A combination product comprising  
(i) a compound according to any one of claims 1 to 20; and  
(ii) another compound effective in the treatment of a respiratory, urological or gastrointestinal disease or disorder  
for simultaneous, separate or sequential use.
- 25 31. A combination product according to claim 30 comprising  
(i) a compound according to any one of claims 1 to 20; and  
(ii) a  $\beta_2$  agonist, steroid, antiallergic drug, phosphodiesterase IV inhibitor and/or leukotriene D4 (LTD4) antagonist for simultaneous, separate or sequential use in the  
30 treatment of a respiratory disease.

## INTERNATIONAL SEARCH REPORT

International Application No.

EP 03/06708

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D453/02 C07D487/08 A61K31/439 A61K31/407 A61P1/00  
 A61P11/00 A61P13/00 //(C07D487/08,209:00,209:00)

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	CA 2 155 320 A (FUJISAWA PHARMACEUTICAL CO) 19 August 1993 (1993-08-19) cited in the application page 1; claim 1 ---	1,2,24, 27
A	WO 01 04118 A (ALMIRALL PRODESFARMA SA) 18 January 2001 (2001-01-18) page 1, line 7 - line 18; claims 1,31,33 ---	1,2,24, 25,27
A	EP 0 930 298 A (BANYU PHARMA CO LTD) 21 July 1999 (1999-07-21) claims 1,15 ---	1,2,24, 25,27
P,A	WO 02 051841 A (ALMIRALL PRODESFARMA ) 4 July 2002 (2002-07-04) claims 1,22,23,25 --- -/--	1,2,25, 27



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

## \* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

1 October 2003

Date of mailing of the international search report

09/10/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
 NL - 2280 HV Rijswijk  
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl  
 Fax: (+31-70) 340-3016

Authorized officer

Alfaro Faus, I

## INTERNATIONAL SEARCH REPORT

International Application No.

/EP 03/06708

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, A	WO 02 053564 A (ALMIRALL PRODESFARMA) 11 July 2002 (2002-07-11) claims 1, 31, 33	1, 2, 25, 27

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No.

EP 03/06708

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
CA 2155320	A	19-08-1993	CA 2155320 A1	19-08-1993
			WO 9316048 A1	19-08-1993
WO 0104118	A	18-01-2001	ES 2165768 A1	16-03-2002
			AT 235492 T	15-04-2003
			AU 6433000 A	30-01-2001
			BG 106301 A	30-08-2002
			BR 0012434 A	02-04-2002
			CA 2381165 A1	18-01-2001
			CN 1373760 T	09-10-2002
			CZ 20020121 A3	13-11-2002
			DE 60001840 D1	30-04-2003
			DK 1200431 T3	21-07-2003
			EE 200200017 A	15-04-2003
			WO 0104118 A2	18-01-2001
			EP 1200431 A2	02-05-2002
			HU 0202100 A2	28-10-2002
			JP 2003504368 T	04-02-2003
			NO 20020180 A	13-03-2002
			SK 432002 A3	01-04-2003
			TR 200200768 T2	22-07-2002
			US 2003055080 A1	20-03-2003
EP 0930298	A	21-07-1999	AT 229941 T	15-01-2003
			AU 716050 B2	17-02-2000
			AU 3635197 A	25-02-1998
			BG 103114 A	30-11-1999
			BR 9711108 A	17-08-1999
			DE 69718026 D1	30-01-2003
			DE 69718026 T2	10-07-2003
			EE 9900038 A	16-08-1999
			EP 0930298 A1	21-07-1999
			JP 3063164 B2	12-07-2000
			NO 990472 A	01-02-1999
			NZ 333842 A	25-05-2001
			SK 12299 A3	16-05-2000
			CN 1226888 A	25-08-1999
			CZ 9900331 A3	14-07-1999
			ES 2188961 T3	01-07-2003
			HR 970426 A1	31-08-1998
			HU 9902381 A2	29-11-1999
			WO 9805641 A1	12-02-1998
			JP 3282617 B2	20-05-2002
			JP 2000178231 A	27-06-2000
			JP 3282618 B2	20-05-2002
			JP 2000169449 A	20-06-2000
			KR 2000022214 A	25-04-2000
			PL 331431 A1	19-07-1999
			TR 9900204 T2	21-01-2000
			TR 200001482 T2	21-11-2000
			US 6040449 A	21-03-2000
			US 5948792 A	07-09-1999
			ZA 9706813 A	11-02-1998
WO 02051841	A	04-07-2002	WO 02051841 A1	04-07-2002
			EP 1345937 A1	24-09-2003
			NO 20032889 A	08-08-2003



# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

EP 03/06708

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 02053564	A	11-07-2002	CA 2433128 A1	11-07-2002
			WO 02053564 A2	11-07-2002
			NO 20033002 A	30-06-2003

---